

History

Introduction

Positive symptoms of schizophrenia, including hallucinations and delusions, have been described throughout ancient Egyptian, Hindu, Chinese, Greek, and Roman writings. Emil Kraepelin, a German physician, was one of the first to classify schizophrenia using the term "dementia praecox", meaning premature dementia. The term "schizophrenia" was coined by Eugen Bleuler around 1910, and roughly translates to "splitting of the mind", a term not meant to suggest a split personality, but to describe a separation of psychological functions: cognition, percept, and affect. Since then, the definition of schizophrenia has continued to change. This summary topic presents the evidence on the history of schizophrenia gained from systematic reviews.

Method

We have included only systematic reviews with detailed literature search, methodology, and inclusion/exclusion criteria that were published in full text, in English, from the year 2000. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. Reviews with pooled data are prioritized for inclusion. Reviews reporting fewer than 50% of items on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses ([PRISMA](#)¹) checklist have been excluded from the library. The evidence was graded guided by the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach². The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found two systematic reviews that met our inclusion criteria^{3, 4}.

- Low quality evidence suggests that the general public in ancient Greece and Rome may have had an awareness of psychotic disorders, but no reference is made to a condition that would meet modern diagnostic criteria for schizophrenia.
- Low quality evidence is unclear about the magnitude of any changes over time in incidence rates of schizophrenia in the UK.

History

Evans K, McGrath J, Milns R

Searching for schizophrenia in ancient Greek and Roman literature: a systematic review

Acta Psychiatrica Scandinavica 2003; 107: 323-330

[View review abstract online](#)

Comparison	Rates of schizophrenia in ancient Greek and Roman literature.
Summary of evidence	Low quality evidence (unable to assess consistency or precision, indirect) is unable to determine the rates of schizophrenia from ancient Greek and Roman literature. However, the literature suggests that the general public may have had an awareness of psychotic disorders, but no reference is made to a condition that would meet modern diagnostic criteria for schizophrenia.
Rates of schizophrenia in ancient Rome and Greece	
The authors report that the general public in ancient Rome and Greece may have had an awareness of psychotic disorders, as psychotic symptoms were described in works of fiction and in historical accounts, however but no reference is made to a condition that would meet modern diagnostic criteria for schizophrenia.	
Consistency in results[‡]	Unable to assess; no measure of consistency is reported.
Precision in results[§]	Unable to assess; no measure of precision is reported.
Directness of results	Indirect assessments.

Kirkbride JB, Errazuriz A, Croudace TJ, Morgan C, Jackson D, Boydell J, Murray RM, Jones PB

Incidence of Schizophrenia and Other Psychoses in England, 1950–2009: A Systematic Review and Meta-Analyses

PLoS One 2012; 7(3): e1660

[View review abstract online](#)

Comparison	Changes over time in incidence of schizophrenia in the UK.
-------------------	---

History

Summary of evidence	Low quality evidence (appears inconsistent, unable to assess precision) is unclear about the magnitude of any changes over time in incidence rates of schizophrenia in the UK.
Changes over time	
<p>Of 9 studies (N = unclear), 3 studies conducted in London reported increased incidence rates from 1965 to 1997. Study authors suggest this may be due to increases in the ethnic immigrant population in the area.</p> <p>2 studies in Nottingham, 1 in Oxfordshire and 1 in England reported decreased incidence rates from 1970 to 1999. Decline matched by corresponding increase in other non-affective psychoses.</p> <p>2 studies in Nottingham reported no changes over time.</p>	
Consistency in results	Unable to assess; no measure of consistency is reported.
Precision in results	Unable to assess; no measure of precision is reported.
Directness of results	Direct

History

Explanation of technical terms

*Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small⁵.

† Different effect measures are reported by different reviews.

Correlation coefficients (eg, r) indicate the strength of association or relationship between variables. They are an indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (*b*) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the dependent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified (100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardized mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. 0.2 represents a small effect, 0.5 a moderate effect, and 0.8 and over represents a large effect⁵.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk

History

factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, an RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. An RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or $< 0.2^6$. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

‡ Inconsistency refers to differing estimates of treatment effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent substantial heterogeneity and 75% to 100%: considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not

weighted mean differences) is considered imprecise if the upper or lower confidence limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, this criteria should be relaxed⁷.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available so is inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.

History

References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Evans K, McGrath J, Milns R (2003): Searching for schizophrenia in ancient Greek and Roman literature: a systematic review. *Acta Psychiatrica Scandinavica* 107: 323-30.
4. Kirkbride JB, Errazuriz A, Croudace TJ, Morgan C, Jackson D, Boydell J, *et al.* (2012): Incidence of schizophrenia and other psychoses in England, 1950-2009: A systematic review and meta-analyses. *PLoS ONE* 7: e31660.
5. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
6. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
7. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. *Version 3.2 for Windows*.