

Incidence in migrants

Introduction

The incidence of schizophrenia refers to how many new cases there are per population in a specified time period. It is different from prevalence, which refers to how many existing cases there are at a particular point in time. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of developing schizophrenia. It takes into account the size of the underlying population sample and its age structure over the duration of observation¹.

Differences in the incidence of a disorder can provide clues to its possible causes. For example, a population register with information gained from consensus data helps to identify all adults who were born within a certain time period (an age cohort) and where they were born. Cross linking this information with a mental health register can be used to identify those who received treatment for schizophrenia over particular times. This can provide information regarding the incidence of schizophrenia within different migrant groups.

The term “migrant” usually refers to first generation migrants - people with a foreign birth place, however some studies also include locally born offspring, or second generation migrants in their analyses. Any association observed between migrant status and increased incidence of schizophrenia has stimulated a great deal of research and explanatory hypotheses, including additional stress relating to migration and settling into a new country, and possible issues with discrimination. Other explanations include a tendency for at-risk individuals to migrate and underlying genetic variances across cultures.

Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2000 that report results separately for people with a diagnosis of schizophrenia, schizoaffective disorder, schizophreniform disorder or first episode schizophrenia. Reviews were identified by searching the databases MEDLINE, EMBASE, CINAHL, Current Contents, PsycINFO and the Cochrane library. Hand searching reference lists of identified reviews was also conducted. When multiple copies of reviews were found, only the most recent version was included. Reviews with pooled data are given priority for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses ([PRISMA](#)) checklist that describes a preferred way to present a meta-analysis². Reviews with less than 50% of items checked have been excluded from the library. The PRISMA flow diagram is a suggested way of providing information about studies included and excluded with reasons for exclusion. Where no flow diagram has been presented by individual reviews, but identified studies have been described in the text, reviews have been checked for this item. Note that early reviews may have been guided by less stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons, imprecise or sparse data and high probability of

Incidence in migrants

reporting bias. It may also be downgraded if risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large, there is a dose dependent response or if results are reasonably consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)³. The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found five systematic reviews that met our inclusion criteria⁴⁻⁸.

- Moderate to high quality evidence finds the incidence rate of schizophrenia is higher in migrants than in native-born populations.
- Moderate quality evidence indicates increased incidence of schizophrenia for both first and second generation migrant populations, particularly for migrants with black skin and those living in the UK, The Netherlands and Scandinavian countries.
- Moderate quality evidence suggests small to medium-sized effects of increased incidence of psychotic disorders in refugee groups after migration (up to 10 years) compared to the native-born population and compared to non-refugee migrants. This risk was highest in men and in refugees from the Middle East.

Incidence in migrants

Bourque F, van der Ven E, Malla A

A meta-analysis of the risk for psychotic disorders among first- and second-generation immigrants

Psychological Medicine 2010; 21: 1-14

[View review abstract online](#)

| | |
|---|--|
| Comparison | Incidence in first and second generation migrants vs. native born populations. |
| Summary of evidence | Moderate quality evidence (large samples, mostly direct, imprecise, inconsistent across and within subgroups) indicates increased incidence of schizophrenia for both first and second generation migrant populations, particularly for migrants with black skin and those living in the UK, The Netherlands and Scandinavian countries. |
| First generation migrants | |
| <p><i>Significant increased incidence of schizophrenia in first generation migrants:</i> 19 population level studies, N = 5508, incidence rate ratio = 2.3, 95%CI = 2.0 to 2.7, $Q_W = 1071, I^2 = 94.4\%, p < 0.01$</p> <p><i>Subgroup analysis – no differences in risk between male and female first generation migrants:</i></p> <p>Males; incidence rate ratio = 2.1, 95%CI = 1.7 to 2.6 $Q_W = 654.2, I^2 = 94.8\%, p < 0.01$</p> <p>Females; incidence rate ratio = 2.4, 95%CI = 1.9 to 2.9 $Q_W = 398.8, I^2 = 91.5\%, p < 0.01$</p> <p>$Q_B = 0.49, NS$</p> | |
| <p><i>Subgroup analysis – risk is highest for first generation migrants where black skin is the majority in their country of origin:</i></p> <p>Black; incidence rate ratio = 4.0, 95%CI = 3.4 to 4.6 $Q_W = 80.8, I^2 = 79\%, p < 0.01$</p> <p>Other; incidence rate ratio = 2.0, 95%CI = 1.6 to 2.5 $Q_W = 97.8, I^2 = 84.7\%, p < 0.01$</p> <p>White; incidence rate ratio = 1.8, 95%CI = 1.6 to 2.1 $Q_W = 175.4, I^2 = 89.7\%, p < 0.01$</p> <p>$Q_B = 57.2, p < 0.01$</p> | |

Incidence in migrants

Subgroup analysis – risk is highest for first generation migrants from black African and black Caribbean ethno-racial groups:

White; incidence rate ratio = 1.8, 95%CI = 1.6 to 2.1

$Q_W = 175.4, I^2 = 89.7\%, p < 0.01$

Black Caribbean; incidence rate ratio = 3.9, 95%CI = 3.4 to 4.6

$Q_W = 42.6, I^2 = 74.2\%, p < 0.01$

Black African; incidence rate ratio = 4.3, 95%CI = 2.8 to 6.8

$Q_W = 38.1, I^2 = 86.9\%, p < 0.01$

Asian; incidence rate ratio = 1.7, 95%CI = 1.3 to 2.3

$Q_W = 32.2, I^2 = 81.4\%, p < 0.01$

Middle East; incidence rate ratio = 2.3, 95%CI = 1.4 to 4.0

$Q_W = 31.9, I^2 = 87.5\%, p < 0.01$

$Q_B = 61.8, p < 0.01$

Subgroup analysis – risk is highest for first generation migrants living in the UK, The Netherlands and Scandinavia:

Israel; incidence rate ratio = 1.5, 95%CI = 1.1 to 2.1

$Q_W = 12.5, I^2 = 51.9\%, \text{NS (trend)}$

The Netherlands; incidence rate ratio = 2.5, 95%CI = 2.0 to 3.2

$Q_W = 61.2, I^2 = 85.3\%, p < 0.01$

Scandinavia; incidence rate ratio = 2.3, 95%CI = 1.9 to 2.7

$Q_W = 185.0, I^2 = 92.4\%, p < 0.01$

UK; incidence rate ratio = 2.8, 95%CI = 2.2 to 3.5

$Q_W = 237.2, I^2 = 93\%, p < 0.01$

$Q_B = 9.5, p < 0.05$

Subgroup analysis – no significant difference in risk based on study setting:

Mixed urban/rural study setting; incidence rate ratio = 2.2, 95%CI = 1.9 to 2.6

$Q_W = 848.0, I^2 = 95.4\%, p < 0.01$

Urban study setting; incidence rate ratio = 2.7, 95%CI = 2.0 to 3.6

$Q_W = 179.1, I^2 = 88.8\%, p < 0.01$

$Q_B = 1.4, \text{NS}$

Subgroup analysis – no significant differences between studies with different sample acquisition methods:

First admission; incidence rate ratio = 2.2, 95%CI = 1.9 to 2.6

Incidence in migrants

| |
|--|
| <p>$Q_W = 961.2, I^2 = 95.5\%, p < 0.01$ First contact; incidence rate ratio = 2.9, 95%CI = 2.1 to 4.0 $Q_W = 103.56, I^2 = 84.5\%, p < 0.01$ $Q_B = 2.2, NS$</p> |
| <p><i>Subgroup analysis – no significant difference in risk based on diagnostic tool:</i> DSM-IV; incidence rate ratio = 2.0, 95%CI = 1.5 to 2.5 $Q_W = 20.4, I^2 = 65.7\%, p < 0.01$ ICD; incidence rate ratio = 2.2, 95%CI = 1.9 to 2.7 $Q_W = 570.2, I^2 = 94.6\%, p < 0.01$ Non-standardized; incidence rate ratio = 2.7, 95%CI = 2.1 to 3.5 $Q_W = 444.4, I^2 = 95.5\%, p < 0.01$ $Q_B = 3.0, NS$</p> |
| <p><i>Subgroup analysis – no significant differences between high and average/low quality studies:</i> High quality studies; incidence rate ratio = 2.2, 95%CI = 1.7 to 2.5 $Q_W = 37.6, I^2 = 65.4\%, p < 0.01$ Average and low quality studies; incidence rate ratio = 2.4, 95%CI = 2.1 to 2.8 $Q_W = 1010.5, I^2 = 95.4\%, p < 0.01$ $Q_B = 1.6, NS$</p> |
| <p>Second generation migrants</p> |
| <p><i>Significant increased incidence of schizophrenia for second generation migrants:</i> 10 population level studies, N = 4422 Incidence rate ratio = 2.1, 95%CI = 1.8 to 2.5 $Q_W = 302, I^2 = 91.1\%, 4.5, p < 0.01$</p> |
| <p><i>Subgroup analysis – no differences in risk between male and female second generation migrants:</i> Males; Incidence rate ratio = 2.5, 95%CI = 1.8 to 3.4 $Q_W = 56.7, I^2 = 78.8\%, p < 0.01$ Females; Incidence rate ratio = 3.0, 95%CI = 2.1 to 4.2 $Q_W = 33.2, I^2 = 63.9\%, p < 0.01$ $Q_B = 0.60, NS$</p> |

Incidence in migrants

Subgroup analysis – highest risk is for second generation migrants where black skin is the majority in their country of origin:

Black; Incidence rate ratio = 5.4, 95%CI = 3.2 to 8.8

$Q_W = 28.4$, $I^2 = 78.9\%$, $p < 0.01$

Other; Incidence rate ratio = 2.0, 95%CI = 1.0 to 4.0

$Q_W = 15.3$, $I^2 = 73.8\%$, $p < 0.01$

White; Incidence rate ratio = 1.9, 95%CI = 1.2 to 3.0

$Q_W = 23.5$, $I^2 = 87.2\%$, $p < 0.01$

$Q_B = 10.6$, $p < 0.01$

Subgroup analysis – risk is highest for second generation migrants from black African and black Caribbean ethno-racial groups:

White; incidence rate ratio = 2.3, 95%CI = 2.1 to 2.7

$Q_W = 1.17$, $I^2 = 0\%$, $p < 0.05$

Black Caribbean; incidence rate ratio = 5.8, 95%CI unclear

$Q_W = 26.3$, $I^2 = 77.2\%$, $p < 0.01$

Black African; incidence rate ratio = 3.7, 95%CI = 2.2 to 6.3

Q_W , I^2 , p , NA – one effect size

Asian; incidence rate ratio = 1.3, 95%CI = 0.8 to 2.1

$Q_W = 0.06$, $I^2 = 0\%$, p unclear

Middle East; incidence rate ratio = 2.3, 95%CI = 1.4 to 4.0

$Q_W = 2.93$, $I^2 = 65.8\%$, $p < 0.01$

$Q_B = 19.9$, $p < 0.01$

Subgroup analysis – risk is highest for second generation migrants living in the UK and The Netherlands:

Israel; incidence rate ratio = 1.1, 95%CI = 0.9 to 1.3

$Q_W = 12.8$, $I^2 = 68.8\%$, $p < 0.05$

The Netherlands; incidence rate ratio = 3.0, 95%CI = 2.1 to 4.4

$Q_W = 7.2$, $I^2 = 30.7\%$, NS

Scandinavia; incidence rate ratio = 1.8, 95%CI = 1.6 to 2.0

$Q_W = 46.0$, $I^2 = 84.8\%$, $p < 0.01$

UK; incidence rate ratio = 3.7, 95%CI = 2.1 to 6.6

$Q_W = 64.5$, $I^2 = 87.6\%$, $p < 0.01$

$Q_B = 34.1$, $p < 0.01$

Incidence in migrants

| | |
|--|---------------------------------------|
| <p><i>Subgroup analysis –no significant difference in risk based on study setting:</i></p> <p>Mixed urban/rural study setting; incidence rate ratio = 1.7, 95%CI = 1.5 to 2.0 $Q_W = 48.2, I^2 = 83.4\%, p < 0.01$</p> <p>Urban study setting; incidence rate ratio = 2.6, 95%CI = 1.7 to 3.9 $Q_W = 253.1, I^2 = 92.9\%, p < 0.01$ $Q_B = 3.0, NS$ (trend)</p> | |
| <p><i>Subgroup analysis – risk is higher in second generation samples from first contact studies:</i></p> <p>First admission; incidence rate ratio = 1.6, 95%CI = 1.3 to 1.8 $Q_W = 149.8, I^2 = 91.3\%, p < 0.01$</p> <p>First contact; incidence rate ratio = 3.2, 95%CI = 2.1 to 4.7 $Q_W = 67.6, I^2 = 80.8\%, p < 0.01$ $Q_B = 10.6, p < 0.01$</p> | |
| <p><i>Subgroup analysis –no significant difference in risk based on diagnostic tool:</i></p> <p>DSM-IV; incidence rate ratio = 2.4, 95%CI = 1.4 to 4.1 $Q_W = 24.9, I^2 = 79.9\%, p < 0.01$</p> <p>ICD; incidence rate ratio = 1.9, 95%CI = 1.6 to 2.2 $Q_W = 230.5, I^2 = 91.8\%, p < 0.01$</p> <p>Non-standardized; incidence rate ratio = 3.7, 95%CI = 0.4 to 31.2 $Q_W = 4.18, I^2 = 76.07\%, p < 0.01$ $Q_B = 1.1, NS$</p> | |
| <p><i>Subgroup analysis – no significant differences between high and average/low quality studies:</i></p> <p>High quality studies; incidence rate ratio = 2.7, 95%CI = 1.9 to 3.7 $Q_W = 32.2, I^2 = 65.8\%, p < 0.01$</p> <p>Average and low quality studies; incidence rate ratio = 1.8, 95%CI = 1.5 to 2.2 $Q_W = 228.7, I^2 = 93.4\%, p < 0.01$ $Q_B = 3.8, NS$ (trend)</p> | |
| Consistency in results[‡] | Rates expected to vary across regions |
| Precision in results[§] | Unable to assess |
| Directness of results | Direct, apart from skin colour. |

Incidence in migrants

Castillejos MC, Martín-Pérez C, Moreno-Küstner B

A systematic review and meta-analysis of the incidence of psychotic disorders: the distribution of rates and the influence of gender, urbanicity, immigration and socio-economic level

Psychological Medicine 2018; 48: 2101–15

[View review abstract online](#)

| | |
|--|---|
| Comparison | Incidence of schizophrenia in migrants vs. native-born populations. |
| Summary of evidence | Moderate quality evidence (large sample, unable to assess consistency, imprecise, direct) suggests the incidence rate of schizophrenia or schizophreniform disorder is higher in migrants than in native-born populations. |
| Diagnosis of schizophrenia or schizophreniform disorder | |
| <i>A significant increased rate of schizophrenia or schizophreniform disorder in migrants; 4 population-based studies, IRR = 3.08, 95%CI 2.04 to 3.67, p < 0.01</i> | |
| Consistency in results | Unable to assess – heterogeneity measure is not reported. |
| Precision in results | Imprecise |
| Directness of results | Direct |

Dapunt J, Kluge U, Heinz A

Risk of psychosis in refugees: a literature review

Translational Psychiatry 2017; 7: e1149

[View review abstract online](#)

| | |
|----------------------------|--|
| Comparison | Incidence of schizophrenia or other non-affective psychotic disorders in refugees vs. native-born populations and non-refugee migrants. |
| Summary of evidence | Moderate quality evidence (large samples, unable to assess consistency, some imprecision, direct) suggests small to |

Incidence in migrants

SCHIZOPHRENIA LIBRARY

| | |
|--|---|
| | <p>medium-sized effects of increased incidence of psychotic disorders in refugee groups after migration (up to 10 years) compared to the native-born population and compared to non-refugee migrants. This risk was highest in men and in refugees from the Middle East.</p> |
| <p>Refugee status</p> | |
| <p style="text-align: center;">1 Swedish cohort study, N = 1,337,790</p> <p>The incidence rate in refugees was 126.4 per 100 000 person years compared to 80.4 in migrant groups and 38.5 in the Swedish-born population.</p> <p><i>A significant, small to medium-sized effect of higher incidence of non-affective psychoses in the refugee group at follow-up (8.9 million person-years) compared to the Swedish-born population;</i></p> <p style="padding-left: 40px;">Adjusted for age and sex: HR = 3.61, 95%CI 2.87 to 4.53, $p < 0.05$</p> <p>Adjusted for disposable income and population density: HR = 2.90, 95%CI 2.31 to 3.64, $p < 0.05$</p> <p><i>A significant, small effect of higher incidence of non-affective psychoses in the refugee group at follow-up (8.9 million person-years) compared to non-refugee migrants;</i></p> <p style="padding-left: 40px;">Adjusted for age and sex: HR = 1.58, 95%CI 1.26 to 1.99, $p < 0.05$</p> <p>Adjusted for disposable income and population density: HR = 1.66, 95%CI 1.32 to 2.09, $p < 0.05$</p> <p>The risk was higher in men than in women, and higher in refugees from the Middle East than other regions.</p> <p style="text-align: center;">1 Canadian cohort study, N = 4,284,694</p> <p>The incidence rate in refugees was 72.8 per 100 000 person years compared to 51.7 in migrant groups and 55.6 in the general population.</p> <p><i>A significant, small effect of higher incidence of psychotic disorders in the refugee group at follow-up (10 years) compared to non-refugee migrants;</i></p> <p style="padding-left: 40px;">IRR = 1.27, 95%CI 1.04 to 1.56, $p < 0.05$</p> <p><i>The effect was not significant when compared to the general population;</i></p> <p style="padding-left: 40px;">IRR = 1.24, 95%CI 0.86 to 1.81, $p > 0.05$</p> <p>The risk was higher in men than in women.</p> <p style="text-align: center;">1 Danish cohort study, N = 145,695</p> <p><i>A significant, small effect of higher incidence of psychotic disorders in the refugee group at follow-up (8 years) compared to Danish-born people;</i></p> <p style="padding-left: 40px;">IRR = 2.03, 95%CI 1.72 to 2.40, $p < 0.05$</p> <p>The risk was higher in men than in women, and higher in refugees from the Middle East than other</p> | |

Incidence in migrants

SCHIZOPHRENIA LIBRARY

| | |
|-------------------------------|--|
| regions. | |
| Consistency in results | Unable to assess; no measure of consistency is reported. |
| Precision in results | Precise, apart from HRs. |
| Directness of results | Direct |

Jongsma HE, Turner C, Kirkbride JB, Jones PB

International incidence of psychotic disorders, 2002-17: a systematic review and meta-analysis

The Lancet Public Health 2019; 4: e229-e44

[View review abstract online](#)

| | |
|---|--|
| Comparison | Incidence of schizophrenia in migrants vs. native-born populations. |
| Summary of evidence | Moderate to high quality evidence (large samples, inconsistent, precise, direct) suggests the incidence rate of schizophrenia is higher in migrants than in native-born populations. |
| Diagnosis of schizophrenia | |
| <i>A significant increased rate of schizophrenia or schizophreniform disorder in migrants; 6 studies (population and other designs), IRR = 1.41, 95%CI 1.15 to 1.75, p < 0.05, I² = 88%</i> | |
| Consistency in results | Inconsistent |
| Precision in results | Precise |
| Directness of results | Direct |

McGrath J, Saha S, Welham J, El Saadi O, MacCauley C, Chant D

A systematic review of the incidence of schizophrenia: the distribution of rates and the influence of sex, urbanicity, migrant status and methodology

BMC Medicine 2004; 2: 13, doi:10.1186/1741-7015-2-13

Incidence in migrants

SCHIZOPHRENIA LIBRARY

[View review abstract online](#)

| | |
|--|--|
| Comparison | Incidence of schizophrenia in migrants vs. native-born populations. |
| Summary of evidence | Moderate quality evidence (large samples, direct, unable to assess precision) suggests that the incidence of schizophrenia is higher in migrants compared to native-born individuals. |
| Incidence of schizophrenia | |
| <p>24 observational studies in total (worldwide), population level data</p> <p><i>Significantly increased incidence of schizophrenia for migrants compared to native-born populations:</i></p> <p>Median rate ratio (10% and 90% quantiles) = 4.6 (1.0 to 12.8)</p> <p>Difference in harmonic means; $F_{1,13} = 51.8$, $p < 0.001$</p> | |
| Consistency in results | Rates expected to vary across regions. |
| Precision in results | Unable to assess quantiles. |
| Directness of results | Direct measure of migration status |

Explanation of acronyms

CI = confidence interval, IRR = incidence rate ratio, p = statistical probability of obtaining that result ($p < 0.05$ generally regarded as significant), Q_B = Q statistic (chi-square) for the test of differences in effect sizes between groups, Q_w = Q statistic (chi-square) for the test of heterogeneity in results across studies, vs. = versus

Incidence in migrants

Explanation of technical terms

* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results, publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small⁹.

† Different effect measures are reported by different reviews.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified - 100% sensitivity = predict all people who are at high risk as developing psychosis and specificity is the proportion of negatives that are correctly identified - 100% specificity = not predicting anyone as being at high risk if they are truly not.

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure

prior to treatment. Standardized mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. 0.2 represents a small effect, 0.5 a moderate effect, and 0.8 and over represents a large effect⁹.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or < 0.2 ¹⁰. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg, r) indicate the strength of association or relationship between variables. They are an indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (b) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other

Incidence in migrants

independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of treatment effect across trials (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may be considerable heterogeneity and over this is considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, this criteria should be relaxed¹¹.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not

available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available so is inferred from available evidence. These inferred treatment effect sized are of lower quality than those gained from head-to-head comparisons of A and B.

Incidence in migrants

References

1. Goldner EM, Hsu L, Waraich P, Somers JM (2002): Prevalence and incidence studies of schizophrenic disorders: a systematic review of the literature. *Canadian Journal of Psychiatry - Revue Canadienne de Psychiatrie* 47: 833-43.
2. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
3. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
4. McGrath J, Saha S, Welham J, El Saadi O, MacCauley C, Chant D (2004): A systematic review of the incidence of schizophrenia: the distribution of rates and the influence of sex, urbanicity, migrant status and methodology. *BMC Medicine* 2: 13.
5. Bourque F, van der Ven E, Malla A (2011): A meta-analysis of the risk for psychotic disorders among first- and second-generation immigrants. *Psychological Medicine* 41: 897-910.
6. Dapunt J, Kluge U, Heinz A (2017): Risk of psychosis in refugees: a literature review. *Translational Psychiatry* 7: e1149.
7. Castillejos MC, Martín-Pérez C, Moreno-Küstner B (2018): A systematic review and meta-analysis of the incidence of psychotic disorders: the distribution of rates and the influence of gender, urbanicity, immigration and socio-economic level. *Psychological Medicine* 48: 2101–15.
8. Jongsma HE, Turner C, Kirkbride JB, Jones PB (2019): International incidence of psychotic disorders, 2002-17: a systematic review and meta-analysis. *The Lancet Public Health* 4: e229-e44.
9. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
10. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
11. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. Version 32 for Windows