



Policy and law

Introduction

Mental health laws in many countries limit involuntary hospital admissions to patients who meet an obligatory dangerousness criterion (ODC) for risk to themselves or others. This policy approach is in use throughout Australia, the USA, and some areas of Canada and Europe. Alternative criteria implemented in the UK and other parts of Canada and Europe allow involuntary treatment in the absence of dangerousness, on the grounds of an assessed need for treatment if the patient is deemed unable to give consent.

Method

We have included only systematic reviews with detailed literature search, methodology, and inclusion/exclusion criteria that were published in full text, in English, from the year 2000. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. Reviews with pooled data are prioritized for inclusion. Reviews reporting fewer than 50% of items on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses ([PRISMA](#)¹) checklist have been excluded from the library. The evidence was graded guided by the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach². The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found one systematic review that met our inclusion criteria³.

- Moderate to high quality evidence indicates that a requirement for patients to satisfy an obligatory 'dangerousness' criterion to allow compulsory treatment may be associated with a longer duration of untreated psychosis in those regions.



Policy and law

Large MM, Nielssen O, Ryan CJ, Hayes R

Mental health laws that require dangerousness for involuntary admission may delay the initial treatment of schizophrenia

Social Psychiatry and Psychiatric Epidemiology 2008; 43(3): 251-256

[View review abstract online](#)

Comparison	Duration of untreated psychosis in countries with and without obligatory dangerousness criterion for involuntary treatment.
Summary of evidence	<p>Moderate to high quality evidence (large samples, unable to assess consistency, precise, direct) indicates longer DUP is associated with a requirement for patients to fill obligatory dangerousness criteria before compulsory admission.</p> <p>Authors conclude that it is likely that at least some of the increase in DUP in regions with an obligatory dangerousness criterion is a direct result of differences in mental health law.</p>
Duration of untreated psychosis (DUP)	
47 studies, N = 5,849	
<p>The mean DUP in samples from regions with an obligatory dangerousness criterion (ODC) was significantly longer than regions without an ODC. Median values were not significantly different.</p> <p>Obligatory dangerousness criterion: Mean 79.5 weeks (95% CI 63.5 to 95.4), weighted mean 77.7 weeks</p> <p>Average median DUP in ODC: 27.5 weeks (95% CI 17.3 to 37.3 weeks), median 28 weeks</p> <p>No- obligatory dangerousness criterion: Mean 55.6 weeks (95% CI 43.4 to 68.8 weeks), weighted mean 55.7 weeks</p> <p>Average median DUP in non-ODC: 19.9 weeks (95% CI 12.9 to 26.9 weeks), median 16 weeks</p>	
Consistency in results	Unable to assess; no measure of consistency is reported.
Precision in results	Precise
Directness of results	Direct

Explanation of acronyms

CI = Confidence Interval, DUP = Duration of Untreated Psychosis, ODC = Obligatory Dangerousness Criterion



Policy and law

Explanation of technical terms

* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small⁴.

† Different effect measures are reported by different reviews.

Correlation coefficients (eg, r) indicate the strength of association or relationship between variables. They are an indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (b) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified (100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomized trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardized mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. 0.2 represents a small effect, 0.5 a medium effect, and 0.8 and over represents a large effect⁴.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For



Policy and law

example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, an RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. An RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or $< 0.2^5$. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, this criteria should be relaxed⁶.

‡ Inconsistency refers to differing estimates of treatment effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent substantial heterogeneity and 75% to 100%: considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available so is inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence



Policy and law

References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Large MM, Niessen O, Ryan CJ, Hayes R (2008): Mental health laws that require dangerousness for involuntary admission may delay the initial treatment of schizophrenia. *Social Psychiatry and Psychiatric Epidemiology* 43: 251-6.
4. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
5. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
6. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. Version 3.2 for Windows