



Art and drama therapies

Introduction

Art and drama therapies are creative therapies generally used as adjuncts to standard care, which encourage the patient to explore alternative methods of communication with a qualified therapist.

Art therapy is defined by the British Association of Art Therapists as “the use of art materials for self-expression and reflection in the presence of a trained art therapist. Clients who are referred to art therapy need not have previous experience or skill in art, the art therapist is not primarily concerned with making an aesthetic or diagnostic assessment of the client’s image. The overall aim of its practitioners is to enable a client to effect change and growth on a personal level through the use of art materials in a safe and facilitating environment”.

Any benefit of art therapy is likely to come from an interaction of the actual production of artworks and the ensuing relationship that develops with the therapist. It has been suggested that the artwork acts as a buffer which reduces the intensity of the relationship between patient and therapist, which may be more productive for schizophrenia patients.

Drama therapy is defined by the British Association of Drama Therapists as “the use of drama and theatre as a therapeutic process. It is a method of working and playing that uses action methods to facilitate creativity, imagination, learning, insight and growth.”

Any benefit of drama therapy may come from the fundamental ‘make-believe’ aspect, which distances participants from the subject matter, allowing them to work with material that may be sensitive to them. Drama therapy may simply involve expressive games and improvisational activities, which are left un-interpreted, while more intense forms of therapy may involve exploration into the content being expressed.

Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2000 that report results separately for people with a diagnosis of schizophrenia, schizoaffective disorder, schizophreniform disorder or first episode schizophrenia. Reviews were identified by searching the databases MEDLINE, EMBASE, CINAHL, Current Contents, PsycINFO and the Cochrane library. Hand searching reference lists of identified reviews was also conducted. When multiple copies of reviews were found, only the most recent version was included. Reviews with pooled data are prioritised for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist that describes a preferred way to present a meta-analysis¹. Reviews rated as having less than 50% of items checked have been excluded from the library. The PRISMA flow diagram is a suggested way of providing information about studies included and excluded with reasons for exclusion. Where no flow diagram has been presented by individual reviews, but identified studies have been described in the text, reviews have been checked for this item. Note that early reviews may have been guided by less stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons,



Art and drama therapies

imprecise or sparse data and high probability of reporting bias. It may also be downgraded if risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large, there is a dose dependent response or if results are reasonably consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)². The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found four systematic reviews that met our inclusion criteria³⁻⁶.

- Moderate quality evidence finds a medium-sized effect of improved positive symptoms, and a large effect of improved negative symptoms, in people with schizophrenia following Chinese calligraphy.
- Moderate to low quality evidence no differences in long-term (1-2 years) symptoms or functioning between art therapy and an activity control or standard care. There may be some benefit of art therapy for symptoms over standard care in the shorter-term (6 weeks).
- Low quality evidence from one very small study is unable to determine the benefits of drama therapy.



Art and drama therapies

SCHIZOPHRENIA LIBRARY

Attard A, Larkin M

Art therapy for people with psychosis: a narrative review of the literature

Lancet Psychiatry 2016; 3: 1067-78

[View review abstract online](#)

Comparison	Art therapy for improving symptoms and functioning in people with schizophrenia vs. controls.
Summary of evidence	Moderate to low quality evidence (small to medium-sized samples, appears inconsistent, imprecise, direct) no differences in long-term (1-2 years) symptoms or functioning between art therapy and an activity control or standard care. There may be some benefit for symptoms over standard care in the shorter-term (6 weeks).
Symptoms and functioning	
<p>1 RCT, N = 417, compared weekly art therapy, activity, or standard care for 1 year and found participants in the activity group had fewer positive symptoms than those in the art therapy group, with no other differences at 2 years follow-up. Authors concluded that art therapy had no clinical advantage and was not more cost-effective than activity.</p> <p>1 RCT, N = 277, compared weekly art therapy with standard care for 1 year and found no significant differences in symptoms between subgroups that had different severities of negative symptoms or different interests in art therapy.</p> <p>1 pilot RCT, N = 58, compared twice weekly art therapy with standard care for 6 weeks and found the art therapy group had a significant decrease in positive symptoms and functioning, and in negative symptoms at follow-up.</p>	
Consistency in results[‡]	Appears inconsistent.
Precision in results[§]	Unable to assess; no measure of consistency is reported.
Directness of results	Direct



Chu KY, Huang CY, Ouyang WC

Does Chinese calligraphy therapy reduce neuropsychiatric symptoms: A systematic review and meta-analysis

BMC Psychiatry 2018; 18 (1)

[View review abstract online](#)

Comparison	Chinese calligraphy for improving symptoms in people with schizophrenia vs. controls (general care, health education or no care). Treatment duration; 8 weeks to 6 months.
Summary of evidence	Moderate quality evidence (small to medium-sized samples, consistent, precise, indirect) finds a medium-sized effect of improved positive symptoms, and a large effect of improved negative symptoms in people with schizophrenia following Chinese calligraphy.
Symptoms	
<p><i>A medium-sized effect of improved positive symptoms in people with schizophrenia;</i> 4 studies, N = 287, SMD = -0.35, 95%CI -0.59, to -0.12, $p = 0.003$, $I^2 = 0\%$</p> <p><i>A large effect of improved negative symptoms in people with schizophrenia;</i> 4 studies, N = 276, SMD = -1.39, 95%CI -1.65, to -1.12, $p < 0.0001$, $I^2 = 15\%$</p>	
Consistency in results	Consistent
Precision in results	Precise
Directness of results	Indirect; mixed control conditions.

Ruddy R, Milnes D

Art therapy for schizophrenia or schizophrenia-like illnesses

Cochrane Database of Systematic Reviews 2005; (4):CD003728

[View review abstract online](#)

Comparison	Art therapy (as defined by the British Association of Art Therapists) administered for 12-20 weeks, with follow up between 6-9 months as an adjunct to standard care vs. standard care.
Summary of evidence	Moderate to low quality evidence (small samples, imprecise, some inconsistency, direct) suggests art therapy may reduce symptom severity and study attrition.
Mental State	
<i>A significant, medium-sized effect favouring art therapy for reducing symptom severity, as measured by Scale for the Assessment of Negative Symptoms (SANS);</i> 1 RCT, N = 73, SMD = -0.58, 95%CI -1.04 to -0.11, $p = 0.02$	
Social functioning	
<i>No significant effect as measured by the Social Functioning Scale (SFS);</i> 1 RCT, N = 70, SMD = 0.34, 95%CI -0.13 to 0.82, $p = 0.15$	
Quality of life	
<i>No significant effect as measured by the Lancashire Quality of Life Profile (PercQoL);</i> 1 RCT, N = 74, SMD = 0.12, 95%CI -0.33 to 0.58, $p = 0.60$	
Leaving the study early	
<i>A significant, medium-sized effect favouring art therapy for reducing attrition in the medium term;</i> 13-26 weeks: 1 study, N = 47, RR = 0.34, 95%CI 0.15 to 0.80, $p = 0.95$ <i>No significant effect in the short or longer term;</i> Up to 12 weeks: 1 study, N = 90, RR = 0.97, 95%CI 0.41 to 2.29, $p = 0.95$ Greater than 26 weeks: 1 study, N = 47, RR = 0.96, 95%CI 0.50 to 1.60, $p = 0.87$	
Consistency in results	Inconsistent for leaving the study early, medium term. Not applicable for all other outcomes – one RCT only.
Precision in results	Imprecise for all outcomes.
Directness of results	Direct



Art and drama therapies

SCHIZOPHRENIA LIBRARY

Ruddy RA, Dent-Brown K

Drama therapy for schizophrenia or schizophrenia-like illnesses

Cochrane Database of Systematic Reviews 2007; (1): CD005378

[View review abstract online](#)

Comparison	Drama therapy (as defined by the British Association of Drama Therapists) administered for 4-22 weeks as an adjunct to medication and inpatient care vs. medication and inpatient care.
Summary of evidence	Low quality evidence (very small sample, imprecise, direct) is unable to determine the benefits of drama therapy for self-esteem.
Self-esteem	
<p><i>A significant, large effect favouring drama therapy for improved self-esteem as measured by the Self-esteem Scale (SES);</i></p> <p style="text-align: center;">1 RCT, N = 24, SMD = 0.97, 95%CI 0.11 to 1.82, <i>p</i> = 0.03</p> <p><i>A significant, large effect favouring drama therapy for inferiority as measured by the Feelings of Inferiority Scale (FIS);</i></p> <p style="text-align: center;">1 RCT, N = 24, SMD = -1.03, 95%CI -1.89 to -0.17, <i>p</i> = 0.02</p>	
Consistency in results	Not applicable, data reported for 1 RCT only.
Precision in results	Imprecise
Directness of results	Direct

Explanation of acronyms

CI = Confidence Interval, FIS = Feelings of Inferiority Scale, I² = the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance), N = number of participants, *p* = statistical probability of obtaining that result (*p* < 0.05 generally regarded as significant), PercQoL = Lancashire Quality of Life Profile, Q = Q statistic (chi-square) for the test of heterogeneity, RCT = randomised controlled trial, RR = risk ratio, SANS = Scale for the Assessment of Negative Symptoms, SAPS = Scale for the Assessment of Positive Symptoms, SES = Self-esteem Scale, SFS = Social Functioning Scale, SMD = standardised mean difference, vs. = versus

Art and drama therapies

Explanation of technical terms

* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small⁷.

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified (100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. 0.2 represents a small effect, 0.5 a medium effect, and 0.8 and over represents a large treatment effect⁷.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, an RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. An RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or < 0.2 ⁸. lnOR stands for logarithmic OR where a lnOR of 0 shows no difference between groups. Hazard ratios



Art and drama therapies

measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg, r) indicate the strength of association or relationship between variables. They are an indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (b) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of treatment effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent substantial heterogeneity and 75% to 100%: considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula⁷;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the

effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, this criteria should be relaxed⁹.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available so is inferred from available evidence. These inferred treatment effect sized are of lower quality than those gained from head-to-head comparisons of A and B.



Art and drama therapies

References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Ruddy R, Milnes D (2005): Art therapy for schizophrenia or schizophrenia-like illnesses. [update of Cochrane Database Syst Rev. 2003;(2):CD003728; PMID: 12804485]. *Cochrane Database of Systematic Reviews*: CD003728.
4. Ruddy RA, Dent-Brown K (2007): Drama therapy for schizophrenia or schizophrenia-like illnesses. *Cochrane Database of Systematic Reviews*: CD005378.
5. Attard A, Larkin M (2016): Art therapy for people with psychosis: a narrative review of the literature. *Lancet Psychiatry* 3: 1067-78.
6. Chu KY, Huang CY, Ouyang WC (2018): Does Chinese calligraphy therapy reduce neuropsychiatric symptoms: A systematic review and meta-analysis. *BMC Psychiatry* 18 (1) (no pagination).
7. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
8. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
9. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. *Version 3.2 for Windows*