



Cognition in family members

Introduction

Cognitive deficits have been reported in people with schizophrenia. Deficits in memory, attention and executive functioning are most commonly reported, with lesser degree of dysfunction in perceptual and language processes. Cognitive deficits are present early in the course of the disorder and are stable over time, and may be heritable.

First-degree relatives of people with schizophrenia may show attenuated signs of cognitive deficits. If cognitive deficits found in people with schizophrenia are also found in their relatives, this may be suggestive of an underlying genetic basis. This is particularly informative in disorders that display complex inheritance patterns such as schizophrenia.

Method

We have included only systematic reviews with detailed literature search, methodology, and inclusion/exclusion criteria that were published in full text, in English, from the year 2000. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. Reviews with pooled data are prioritized for inclusion. Reviews reporting fewer than 50% of items on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses ([PRISMA](#)¹) checklist have been excluded from the library. The evidence was graded guided by the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach². The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found 12 reviews that met inclusion criteria³⁻¹⁴.

- High quality evidence shows small to medium-sized effects of poor executive functioning (including tasks of attention and language), poor visual memory, verbal memory, short-term and long-term episodic memory, and social cognition (Theory of Mind and negative facial expression recognition) in relatives compared with controls.
- Moderate to high quality evidence suggests small to medium-sized effects of lower IQ, working memory, prospective memory, and slower processing speed, and moderate quality evidence suggests poorer psychomotor and visuospatial ability in relatives compared with controls.
- Compared to relatives of people with bipolar disorder, moderate to high quality evidence suggests relatives of people with schizophrenia show small to medium-sized effects of poorer performance on IQ, verbal memory, working memory, processing speed, verbal fluency, and accuracy on executive functioning tasks.



Bora E, Pantelis C

Theory of mind impairments in first-episode psychosis, individuals at ultra-high risk for psychosis and in first-degree relatives of schizophrenia: Systematic review and meta-analysis

Schizophrenia Research 2013; 144(1-3): 31-36

[View review abstract online](#)

Comparison	Assessing performance on Theory of Mind tasks in relatives of patients with a diagnosis of schizophrenia spectrum disorder vs. healthy controls.
Summary of evidence	High quality evidence (medium to large samples, precise, consistent, direct) suggests a small to medium-sized Theory of Mind impairment in relatives of people with schizophrenia. Lower education in patients contributed to lower Theory of Mind scores.
Combined Theory of Mind score	
<p><i>Small to medium effect showing impaired performance in relatives vs. controls;</i> 12 studies, N = 3,117, $d = 0.37$, 95%CI 0.19 to 0.54, $p < 0.001$, $I^2 = 0\%$</p> <p>Longer duration of education in the control groups vs. relatives explained some of the between-group differences ($B = 0.41$, $p = 0.002$).</p>	
Theory of Mind verbal	
<p><i>Small effect showing impaired performance in relatives vs. controls;</i> 8 studies, N = 2,946, $d = 0.24$, 95%CI 0.13 to 0.33, $p < 0.001$, $I^2 = 0\%$</p>	
Theory of Mind visual	
<p><i>Small to medium effect showing impaired performance in relatives vs. controls;</i> 9 studies, N = 555, $d = 0.36$, 95%CI 0.10 to 0.63, $p < 0.001$, $I^2 = 0\%$</p>	
Eyes task	
<p><i>No differences between groups;</i> 5 studies, N = 261, $d = 0.19$, 95%CI -0.10 to 0.48, $p = 0.19$, $I^2 = 0\%$</p>	



Cognition in family members

Consistency[‡]	Consistent
Precision[§]	Precise
Directness	Direct

Bora E

A comparative meta-analysis of neurocognition in first-degree relatives of patients with schizophrenia and bipolar disorder

European Psychiatry: the Journal of the Association of European Psychiatrists 2017; 45: 121-8

[View review abstract online](#)

Comparison 1	Cognition in first-degree relatives of people with schizophrenia vs. controls.
Summary of evidence	Moderate to high quality evidence (large samples, mostly precise, some inconsistency, direct) suggests small to medium-sized effects of poorer performance on IQ, verbal, working and visual memory, processing speed, sustained attention, executive functioning and verbal fluency in relatives of people with schizophrenia.
IQ	
<i>Significant, medium-sized effect of lower IQ in relatives of schizophrenia patients; 13 studies, N = 1,618, d = 0.48, 95CI% 0.29 to 0.67, p < 0.001, I² = 67%, p < 0.001</i>	
Memory	
<i>Significant, medium-sized effects of poorer memory in relatives of schizophrenia patients;</i> Verbal memory: 7 studies, N = 965, d = 0.46, 95CI% 0.23 to 0.69, p < 0.001, I ² = 61%, p = 0.02 Working memory: 9 studies, N = 793, d = 0.57, 95CI% 0.41 to 0.77, p < 0.001, I ² = 21%, p = 0.25 Visual memory (trend): 4 studies, N = 464, d = 0.71, 95CI% -0.02 to 1.44, p = 0.06, I ² = 94%, p < 0.001	
Processing speed	



Cognition in family members

<p><i>Significant, medium-sized effects of slower processing speed in relatives of schizophrenia patients;</i> 8 studies, N = 764, $d = 0.59$, 95CI% 0.29 to 0.88, $p < 0.01$, $I^2 = 76%$, $p < 0.001$</p>	
<p>Attention</p>	
<p><i>Significant, small effect of poorer sustained attention in relatives of schizophrenia patients;</i> 8 studies, N = 1,835, $d = 0.20$, 95CI% 0.11 to 0.29, $p < 0.001$, $I^2 = 0%$, $p = 0.86$</p>	
<p>Executive functioning</p>	
<p><i>Significant, medium-sized effects of poorer executive functioning in relatives of schizophrenia patients;</i> Executive functioning: 11 studies, N = 964, $d = 0.58$, 95CI% 0.32 to 0.85, $p < 0.001$, $I^2 = 76%$ $p < 0.001$ Accuracy: 9 studies, N = 706, $d = 0.43$, 95CI% 0.25 to 0.60, $p < 0.001$, $I^2 = 24%$ $p = 0.24$ Speed: 6 studies, N = 545, $d = 0.58$, 95CI% 0.20 to 0.96, $p = 0.003$, $I^2 = 76%$ $p < 0.001$</p>	
<p>Language</p>	
<p><i>Significant, medium-sized effect of poorer verbal fluency in relatives of schizophrenia patients;</i> 7 studies, N = 536, $d = 0.56$, 95CI% 0.36 to 0.76, $p < 0.001$, $I^2 = 24%$, $p = 0.24$</p>	
Consistency	Inconsistent for IQ, visual memory, processing speed, and executive functioning.
Precision	Precise, apart from visual memory.
Directness	Direct
Comparison 2	Cognition in first-degree relatives of people with schizophrenia vs. first-degree relatives of people with bipolar disorder.
Summary of evidence	Moderate to high quality evidence (large samples, precise, some inconsistency, direct) suggests small to medium-sized effects of poorer performance on IQ, verbal memory, working memory, processing speed, verbal fluency, and accuracy of executive functioning tasks in relatives of people with schizophrenia. There were no significant differences in speeded executive functioning tasks, IQ, visual memory, and sustained attention.
<p>IQ</p>	



Cognition in family members

<p><i>Significant, small to medium-sized effect of lower IQ in relatives of schizophrenia patients;</i> 13 studies, N = 1,263, $d = 0.38$, 95%CI 0.14 to 0.62, $p < 0.001$, $I^2 = 72%$, $p < 0.01$ Shorter duration of education in the relatives of people with schizophrenia was associated with greater between-group differences in global cognition.</p>	
Memory	
<p><i>Significant, small to medium-sized effects of poorer memory in relatives of schizophrenia patients;</i> Verbal memory: 8 studies, N = 815, $d = 0.28$, 95%CI 0.04 to 0.53, $p = 0.02$, $I^2 = 49%$, $p = 0.06$ Working memory: 10 studies, N = 589, $d = 0.42$, 95%CI 0.18 to 0.66, $p < 0.001$, $I^2 = 59%$, $p = 0.009$ There were no differences in visual memory.</p>	
Processing speed	
<p><i>Significant, small effect of slower processing speed in relatives of schizophrenia patients;</i> 9 studies, N = 699, $d = 0.30$, 95%CI 0.06 to 0.53, $p = 0.01$, $I^2 = 56%$, $p = 0.02$</p>	
Executive functioning	
<p><i>Significant, small effect of poorer executive functioning in relatives of schizophrenia patients;</i> Accuracy: 10 studies, N = 753, $d = 0.27$, 95%CI 0.12 to 0.42, $p < 0.001$, $I^2 = 4%$, $p = 0.40$ There were no significant differences on speeded tasks.</p>	
Language	
<p><i>Significant, small effect of poorer verbal fluency in relatives of schizophrenia patients;</i> 7 studies, N = 431, $d = 0.24$, 95%CI 0 to 0.47, $p = 0.05$, $I^2 = 33%$, $p = 0.18$</p>	
Consistency	Inconsistent for IQ, verbal memory, working memory and processing speed.
Precision	Precise
Directness	Direct

Cohen AS, Brown LA, Auster TL

Olfaction, “olfiction,” and the schizophrenia-spectrum: An updated meta-analysis on identification and acuity



Schizophrenia Research 2012; 135: 152-157 View review abstract online	
Comparison	Olfactory identification and acuity in people with a family history of schizophrenia.
Summary of evidence	Moderate quality evidence (precise, direct, unable to assess consistency) suggests no impairment in olfactory performance.
Olfactory performance	
<i>No significant differences were reported;</i> 9 studies, N = 517, $d = -0.21$, 95%CI -0.53 to 0.12, $p > 0.05$	
Consistency	Unable to assess
Precision	Precise
Directness	Direct

<i>Dickinson D, Ramsey ME, Gold JM</i>	
Overlooking the Obvious: A meta-analytic comparison of digit symbol coding tasks and other cognitive measures in schizophrenia	
Archives of General Psychiatry 2007; 64: 532-542 View review abstract online	
Comparison	Cognitive functioning in relatives of people with schizophrenia vs. controls.
Summary of evidence	Moderate to high quality evidence (medium-sized samples, unable to assess consistency, precise, direct) suggests a medium effect of poorer processing speed, and a small to medium effect of poorer sustained attention, episodic memory and executive functioning in relatives compared with controls. There were no differences in IQ, working memory, or letter fluency.
Attention	
<i>A small effect suggests relatives showed poorer sustained attention on the degraded stimulus CPT</i>	



Cognition in family members

<i>compared to controls;</i> 3 studies, N = 171, $g = -0.39$, 95%CI -0.70 to -0.08, $p < 0.05$, Q p not reported	
IQ	
<i>No difference IQ;</i> 4 studies, N = 315, $g = -0.20$, 95%CI -0.43 to 0.04, $p > 0.05$, Q p not reported	
Memory	
<i>A small to medium effect suggests relatives showed poorer episodic memory performance on the word list learning task compared to controls;</i> 3 studies, N = 193, $g = -0.45$, 95%CI -0.72 to -0.13, $p < 0.05$, Q p not reported <i>No difference in working memory performance on the digit span total;</i> 4 studies, N = 235, $g = -0.23$, 95%CI -0.50 to 0.03, $p > 0.05$, Q p not reported	
Language	
<i>No difference in fluency performance on the letter fluency task;</i> 3 studies, N = 224, $g = -0.22$, 95%CI -0.49 to 0.05, $p > 0.05$, Q p not reported	
Processing speed	
<i>Medium effect sizes suggest relatives showed poorer processing speed performance compared to controls on;</i> Digit symbol coding: 7 studies, N = 504, $g = -0.62$, 95%CI -0.80 to -0.43, $p < 0.05$ TMT-A: 3 studies, N = 193, $g = -0.51$, 95%CI -0.80 to -0.21, $p < 0.05$, Q p not reported	
Executive functioning	
<i>A small effect size suggests relatives showed poorer executive functioning on the WCST categories compared to controls;</i> 4 studies, N = 269, $g = -0.26$, 95%CI -0.50 to 0.00, $p < 0.05$, Q p not reported <i>A medium effect size suggests relatives showed poorer executive functioning on the trail making test part B compared to controls;</i> 3 studies, N = 193, $g = -0.54$, 95%CI -0.82 to -0.23, $p < 0.05$, Q p not reported	
Consistency in results	Unable to assess; no measure of consistency is reported.
Precision in results	Precise
Directness of results	Direct



Jameson KG, Nasrallah HA, Northern TG, Welge JA

Executive function impairment in first-degree relatives of persons with schizophrenia: A meta-analysis of controlled studies

Asian Journal of Psychiatry 2011; 4: 96

[View review abstract online](#)

Comparison	Executive functioning in first-degree relatives of people with schizophrenia vs. controls.
Summary of evidence	Moderate quality evidence (large samples, inconsistent or imprecise, direct, potential publication bias) suggests poorer executive functioning in first-degree relatives of people with schizophrenia.
Executive functioning	
<p><i>A small to medium effect suggests significantly fewer categories achieved and significantly more perseverative errors on the WCST in relatives of people with schizophrenia compared to controls;</i></p> <p>Categories achieved: 17 studies, N = 1,602, $d = -0.34$, 95%CI -0.50 to 0.18, p value not reported</p> <p>Authors report no heterogeneity ($p = 0.273$), but potential publication bias ($p = 0.08$)</p> <p>Perseverative errors: 23 studies, N = 2,173, $d = 0.26$, 95%CI 0.06 to 0.46, p value not reported</p> <p>Authors report significant heterogeneity ($p < 0.001$) and potential publication bias ($p = 0.06$)</p>	
Consistency	Consistent for categories, inconsistent for perseverative errors.
Precision	Precise for perseverative errors, imprecise for categories.
Directness	Direct

Lin SZ, Wu YK, Su YA, Si TM

Prospective memory in non-psychotic first-degree relatives of patients with schizophrenia: A meta-analysis

Neuropsychiatric Disease and Treatment 2019; 15: 1563-71

[View review abstract online](#)



Comparison	Prospective memory in first-degree relatives of people with schizophrenia vs. controls.
Summary of evidence	Moderate to high quality evidence (small to medium-sized sample, consistent, precise, direct) suggests poorer facial emotion recognition, particularly negative expressions, in first-degree relatives of people with schizophrenia.
Prospective memory	
<p><i>First-degree relatives of patients with schizophrenia showed impairments in;</i></p> <p>Overall prospective memory: 2 studies, N = 127, SMD = -0.46, 95%CI -0.82 to -0.11, $p = 0.01$, $I^2 = 0\%$</p> <p>Event-based prospective memory: 4 studies, N = 268, SMD = -0.56, 95%CI -0.80 to -0.31, $p < 0.00001$, $I^2 = 0\%$</p> <p>Time-based prospective memory: 4 studies, N = 268, SMD = -0.66, 95%CI -0.90 to -0.41, $p < 0.00001$, $I^2 = 0\%$</p>	
Consistency	Consistent
Precision	Precise
Directness	Direct

Martin D, Croft J, Pitt A, Strelchuk D, Sullivan S, Zammit S

Systematic review and meta-analysis of the relationship between genetic risk for schizophrenia and facial emotion recognition

Schizophrenia Research 2020; 218: 7-13

[View review abstract online](#)

Comparison	Facial emotion recognition in first-degree relatives of people with schizophrenia vs. controls.
Summary of evidence	Moderate to high quality evidence (large sample, inconsistent, precise, direct) suggests poorer facial emotion recognition, particularly negative expressions, in first-degree relatives of people with schizophrenia.
Facial expression recognition	



Cognition in family members

<p><i>A medium-sized effect of poorer facial emotion recognition in first-degree relatives of people with schizophrenia;</i></p> <p>23 studies, N = 2,237, SMD = 0.38 95%CI 0.25 to 0.51, $p < 0.001$, $I^2 = 47%$, $p < 0.007$</p> <p>There was a small, significant effect of poorer recognition of negative valence facial expressions but not positive valence facial expressions.</p>	
Consistency	Consistent for categories, inconsistent for perseverative errors.
Precision	Precise for perseverative errors, imprecise for categories.
Directness	Direct

<p><i>Sitskoorn M, Aleman A, Ebishe S, Appels M, Kahn R</i></p> <p>Cognitive deficits in relatives of patients with schizophrenia: a meta-analysis</p> <p>Schizophrenia Research 2004; 17: 285-295</p> <p>View review abstract online</p>	
Comparison	Cognitive functioning in first-degree relatives of people with schizophrenia vs. healthy controls.
Summary of evidence	High quality evidence (large samples, consistent, precise, direct) finds a small to medium effect of poorer executive functioning as measured by TMT-B, attention as measured by Stroop and TMT-A, and visual and verbal memory in first-degree relatives. Moderate to high quality evidence (inconsistent) suggests poorer performance on executive functioning as measured by WCST, attention as measured by CPT, and verbal fluency.
Executive functioning	
<p><i>Small to medium effect sizes suggest first-degree relatives performed significantly worse on;</i></p> <p>TMT-B: 12 studies, N = 1,424, $d = 0.51$, 95%CI 0.36 to 0.67, $p < 0.0001$, $Q = 12.9$, $p > 0.05$</p> <p>WCST: 19 studies, N = 860, $d = 0.29$ 95%CI 0.14 to 0.43 $p < 0.0001$, $Q = 33.2$, $p < 0.05$</p>	
Attention	
<p><i>Small effect sizes suggest first degree relatives performed significantly worse on;</i></p>	



Cognition in family members

<p>CPT: 11 studies, N = 951, $d = 0.33$, 95%CI 0.09 to 0.57, $p = 0.006$, $Q = 27.6$, $p < 0.01$ Stroop: 8 studies, N = 1,689, $d = 0.28$, 95%CI 0.06 to 0.50, $p = 0.01$, $Q = 11.8$, $p > 0.05$ TMT-A: 10 studies, N = 843, $d = 0.38$, 95%CI 0.23 to 0.53, $p < 0.0001$, $Q = 9.3$, $p > 0.05$</p>	
Language	
<p><i>A small effect size suggest first degree relatives performed significantly worse on verbal fluency;</i> 13 studies, N = 887, $d = 0.35$, 95%CI 0.14 to 0.56, $p = 0.001$, $Q = 30.6$, $p < 0.01$</p>	
Memory	
<p><i>Small to medium effect sizes suggest first-degree relatives performed significantly worse on;</i> CVLT/WMS verbal memory: 15 studies, N = 997, $d = 0.54$, 95%CI 0.43 to 0.66, $p < 0.0001$, $Q = 12.3$, $p > 0.05$ WMS visual reproduction: 8 studies, N = 1,148, $d = 0.30$, 95%CI 0.10 to 0.50, $p = 0.003$, $Q = 11.2$, $p > 0.05$ Digit Span: 10 studies, N = 630, $d = 0.35$, 95%CI 0.19 to 0.50, $p < 0.0001$, $Q = 4.4$, $p > 0.05$</p>	
Consistency	Consistent, apart from language, CPT and WCST.
Precision	Precise
Directness	Direct

Snitz B, MacDonald III A, Carter C

Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: A meta-analytic review of putative endophenotypes

Schizophrenia Bulletin 2006; 32(1): 179-194

[View review abstract online](#)

Comparison	Cognitive functioning in first-degree relatives of people with schizophrenia vs. healthy controls.
Summary of evidence	Moderate quality evidence (large sample, unable to assess consistency or precision, direct) suggests a small to medium effect of poorer executive functioning, verbal and visual memory, psychomotor ability and IQ, and poorer performance on some attention, language and visuospatial tasks in relatives.



Executive functioning

Small to medium effect sizes suggest first-degree relatives performed significantly worse on;

Wisconsin Card Sorting Task (categories): 17 studies, N = 1,114, $g = 0.38$, $p < 0.05$

Wisconsin Card Sorting Task (total errors): 8 studies, N = 635, $g = 0.39$, $p < 0.05$

Wisconsin Card Sorting Task (perseverative errors/responses): 19 studies, N = 1,572, $g = 0.40$, $p < 0.05$

Attention

Small to medium effects sizes suggest first degree relatives performed significantly worse on;

Spatial delayed response (accuracy): 4 studies, N = 236, $g = 0.55$, $p < 0.05$

CPT-AX/ -IP false alarms: 5 studies, N = 450, $g = 0.54$, $p < 0.05$

CPT-AX/ -IP d prime: 8 studies, N = 805, $g = 0.53$, $p < 0.05$

Stroop test- colour naming: 3 studies, N = 200, $g = 0.44$, $p < 0.05$

CPT-X d prime: 8 studies, N = 687, $g = 0.43$, $p < 0.05$

Trails B (time): 16 studies, N = 1,364, $g = 0.41$, $p < 0.05$

Visual cancellation test (accuracy): 4 studies, N = 418, $g = 0.39$, $p < 0.05$

Stroop test- colour-word naming: 5 studies, N = 326, $g = 0.33$, $p < 0.05$

CPT-X hits/omission errors: 3 studies, N = 240, $g = 0.33$, $p > 0.05$

Trails A (time): 11 studies, N = 912, $g = 0.31$, $p < 0.05$

Antisaccade (percentage errors): 6 studies, N = 599, $g = 0.25$, $p < 0.05$

Span of apprehension (accuracy): 6 studies, N = 419, $g = 0.23$, $p < 0.05$

Antisaccade (reaction time): 6 studies, N = 599, $g = 0.19$, $p < 0.05$

There were no significant differences between groups on:

Prosaccade (reaction time): 6 studies, N = 599, $g = -0.00$, $p > 0.05$

Stroop test- word reading: 4 studies, N = 284, $g = 0.07$, $p > 0.05$

Spatial delayed response task (reaction time): 4 studies, N = 236, $g = 0.24$, $p > 0.05$

Spatial span forward: 3 studies, N = 251, $g = 0.25$, $p > 0.05$

CPT-AX/-IP hits/omission errors: 5 studies, N = 450, $g = 0.21$, $p > 0.05$

CPT-X false alarms: 3 studies, N = 240, $g = 0.25$, $p > 0.05$

IQ

A small effect size suggests first-degree relatives performed significantly worse on;

IQ: 9 studies, N = 717, $g = 0.31$, $p < 0.05$



Language

Small to medium effect sizes suggest first-degree relatives performed significantly worse on;

NART/ Wide Range Achievement Test (revised): 6 studies, N = 477, $g = 0.50$, $p < 0.05$

Category fluency tasks: 6 studies, N = 383, $g = 0.68$, $p < 0.05$

Wechsler Adult Intelligence Scale (revised) Vocabulary: 6 studies, N = 604, $g = 0.21$, $p < 0.05$

Letter fluency tasks: 7 studies, N = 533, $g = 0.48$, $p < 0.05$

There was no significant difference between groups in:

WAIS (revised) Information: 3 studies, N = 194 (82 relatives, 112 controls), $g = 0.27$, $p > 0.05$

Memory

Small to medium effect sizes suggest first-degree relatives performed significantly worse on;

Auditory verbal learning task: 3 studies, N = 303, $g = 0.56$, $p < 0.05$

Wechsler Memory Scale – logical memory I: 8 studies, N = 727, $g = 0.49$, $p < 0.05$

Wechsler Memory Scale – logical memory II: 8 studies, N = 848, $g = 0.32$, $p < 0.05$

Wechsler Memory Scale – verbal paired associations: 4 studies, N = 412, $g = 0.42$, $p < 0.05$

Wechsler Memory Scale – visual reproduction I: 7 studies, N = 680, $g = 0.38$, $p < 0.05$

Wechsler Memory Scale – visual reproduction II: 8 studies, N = 908, $g = 0.34$, $p < 0.05$

Digit span forward: 11 studies, N = 983, $g = 0.29$, $p < 0.05$

Digit span backwards: 9 studies, N = 822, $g = 0.27$, $p < 0.05$

Psychomotor

Small effect sizes suggest first-degree relatives performed significantly worse on;

Pegboard task – non-dominant hand: 4 studies, N = 555, $g = 0.26$, $p < 0.05$

Finger tapping – dominant hand: 3 studies, N = 321, $g = 0.33$, $p < 0.05$

Finger tapping – non-dominant hand: 3 studies, N = 321, $g = 0.25$, $p < 0.05$

Pegboard tasks – dominant hand: 4 studies, N = 555, $g = 0.18$, $p < 0.05$

Visuospatial

Small to medium effect sizes suggest first-degree relatives performed significantly worse on;

Design and copy task: 4 studies, N = 329, $g = 0.63$, $p < 0.05$

WAIS-R block design: 7 studies, N = 736, $g = 0.34$, $p < 0.05$

No significant difference in;



Line orientation: 3 studies, N = 353, $g = 0.09$, $p > 0.05$

Consistency	Unable to assess; no measure of consistency is reported.
Precision	Unable to assess; no measure of precision is reported.
Directness	Direct

Szöke A, Schurhoff F, Mathieu F, Meary A, Ionescu S, Leboyer M

Tests of executive functions in first-degree relatives of schizophrenic patients: a meta-analysis

Psychological Medicine 2005; 35: 771-782

[View review abstract online](#)

Comparison	Executive functioning in first-degree relatives of people with schizophrenia vs. healthy controls.
Summary of evidence	High quality evidence (large samples, consistent, precise, direct) shows a small to medium effect of poorer executive functioning in general (also including measures of attention and language), and a large effect of poorer semantic fluency in relatives compared with controls.
Executive functioning Includes attention and language tasks	
<p><i>Small to medium effect sizes suggest first-degree relatives performed significantly worse on;</i></p> <p>WCST (categories): 13 studies, N = 1,619, $g = 0.31$, 95%CI 0.21 to 0.42, $p < 0.05$, $Q p = 0.10$</p> <p>WCST (perseverative): 12 studies, N = 1,261, $g = 0.26$, 95%CI 0.14 to 0.38, $p < 0.05$, $Q p > 0.05$</p> <p>TMT-B: 11 studies, N = 1,179, $g = 0.49$, 95%CI 0.37 to 0.62, $p < 0.05$, $Q p = 0.47$</p> <p>Stroop: 6 studies, N = 563, $g = 0.38$, 95%CI 0.21 to 0.55, $p < 0.05$, $Q p = 0.15$</p> <p>Phonological fluency: 8 studies, N = 664, $g = 0.65$, 95%CI 0.48 to 0.82, $p < 0.05$, $Q p = 0.12$</p> <p style="text-align: center;"><i>With a large effect on:</i></p> <p>Semantic fluency: 5 studies, N = 336, $g = 0.87$, 95%CI 0.64 to 1.10, $p < 0.05$, $Q p = 0.83$</p>	
Consistency	Consistent
Precision	Precise



Directness	Direct
-------------------	--------

Trandafir A, Meary A, Schurhoff F, Leboyer M, Szoke A

Memory tests in adult relatives of schizophrenic patients: a meta-analysis

Schizophrenia Research 2006; 81: 217-226

[View review abstract online](#)

Comparison	Memory in first-degree relatives of people with schizophrenia vs. healthy controls.
Summary of evidence	High quality evidence (medium to large samples, consistent, precise, direct) shows a small to medium effect of poorer visual and verbal memory in relatives compared with controls.

Memory

Small to medium effect sizes suggest first-degree relatives performed significantly worse on;

Verbal paired associates: 4 studies, N = 369, $g = 0.54$, 95%CI 0.33 to 0.75, $p < 0.05$, $Qp = 0.85$

Digit Span forward: 10 studies, N = 748, $g = 0.45$, 95%CI 0.30 to 0.60, $p < 0.05$, $Qp = 0.42$

Digit Span backward: 10 studies, N = 773, $g = 0.35$, 95%CI 0.20 to 0.50, $p < 0.05$, $Qp = 0.14$

Logical Stories immediate: 10 studies, N = 994, $g = 0.47$, 95%CI 0.33 to 0.60, $p < 0.05$, $Qp = 0.20$

Logical Stories delayed (raw): 6 studies, N = 621, $g = 0.38$, 95%CI 0.20 to 0.55, $p < 0.05$, $Qp = 0.26$

Logical Stories delayed (retained): 4 studies, N = 351, $g = 0.18$, 95%CI 0.03 to 0.40, $p < 0.05$, $Qp = 0.52$

Visual reproduction immediate: 8 studies, N = 1,094, $g = 0.17$, 95%CI 0.02 to 0.32, $p < 0.05$, $Qp = 0.18$

Visual reproduction delayed (raw): 4 studies, N = 551, $g = 0.24$, 95%CI 0.05 to 0.33, $p < 0.05$, $Qp = 0.16$

Visual reproduction delayed (retained): 4 studies, N = 351, $g = 0.16$, 95%CI 0.05 to 0.37, $p < 0.05$, $Qp > 0.05$

CVLT: 5 studies, N = 548, $g = 0.30$, 95%CI 0.10 to 0.48, $p < 0.05$, $Qp = 0.65$

Consistency	Consistent
Precision	Precise



Directness	Direct
-------------------	--------

Whyte M, McIntosh A, Johnstone E, Lawrie S

Declarative memory in unaffected adult relatives of patients with schizophrenia: A systematic review and meta-analysis

Schizophrenia Research 2005; 78: 13–26

[View review abstract online](#)

Comparison	Memory in first-degree relatives of people with schizophrenia vs. healthy controls.
Summary of evidence	High quality evidence (medium to large samples, consistent, precise, direct) shows a small to medium effect of poorer short and long term episodic memory in relatives compared with controls. Moderate to high quality evidence (inconsistent) also suggests a small to medium effect of poorer IQ.
Memory	
<p><i>Small to medium effect sizes suggest first-degree relatives performed significantly worse on;</i></p> <p style="text-align: center;"><u>Short-term episodic encoding and retrieval</u></p> <p>Trial 1 list recall: 3 studies, N = 267, $d = 0.65$, 95%CI 0.36 to 0.95, $p < 0.001$, $Qp = 0.99$ Immediate story recall: 10 studies, N = 1,248, $d = 0.53$, 95%CI 0.40 to 0.67, $p < 0.001$, $Qp = 0.59$ Immediate visual recall: 7 studies, N = 941, $d = 0.32$, 95%CI 0.08 to 0.56, $p = 0.008$, $Qp = 0.038$, $I^2 = 54.9$</p> <p style="text-align: center;"><u>Long-term episodic encoding and retrieval</u></p> <p>Delayed story recall: 7 studies, N = 955, $d = 0.52$, 95%CI 0.36 to 0.69 $p < 0.001$, $Qp = 0.64$ CVLT 1–5 total: 6 studies, N = 683, $d = 0.44$, 95%CI 0.25 to 0.63, $p < 0.001$, $Qp = 0.30$ Paired associates: 4 studies, N = 416, $d = 0.41$, 95%CI 0.19 to 0.62, $p < 0.001$, $Qp = 0.50$ Delayed visual recall: 6 studies, N = 867, $d = 0.32$, 95%CI 0.12 to 0.52, $p = 0.002$, $Qp = 0.22$</p> <p style="text-align: center;"><u>Semantic retrieval</u></p> <p>Verbal letter fluency: 12 studies, N = 1,037, $d = 0.42$, 95%CI 0.24 to 0.60, $p < 0.001$, $Qp = 0.40$ Verbal category fluency: 9 studies, N = 901, $d = 0.39$, 95%CI 0.19 to 0.59, $p < 0.001$, $Qp = 0.09$</p>	
IQ	



Cognition in family members

<p><i>Small to medium effect sizes suggest first-degree relatives performed significantly worse on;</i> NART/ Wide Range Achievement Test: 6 studies, N = 850, $d = 0.53$, 95%CI 0.09 to 0.96, $p = 0.017$, $Qp < 0.001$, $I^2 = 81.0$ WAIS-R IQ: 10 studies, N = 1,141, $d = 0.34$, 95%CI 0.07 to 0.61, $p = 0.014$, $Qp < 0.001$, $I^2 = 72.6$</p>	
Consistency	Inconsistent for IQ and visual recall.
Precision	Precise
Directness	Direct

Explanation of acronyms

CI = confidence interval, CPT = Continuous Performance Test, CVLT = California Verbal Learning Test, d = Cohen’s d and g = Hedges’ g = standardised mean differences (see below for interpretation of effect size), I^2 = the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance), IQ = intelligence quotient, N = number of participants, p = statistical probability of obtaining that result ($p < 0.05$ generally regarded as significant), Q = Q statistic for the test of heterogeneity, Q_w = test for within group differences (heterogeneity in study results within a group of studies – measure of study consistency), Q_B = test for between group differences (heterogeneity between groups of studies for an outcome of interest), SMD = standardised mean difference, TMT = Trail Making Test, vs. = versus, WAIS = Wechsler Adult Intelligence Scale, WCST= Wisconsin Card Sorting Test, WMS = Wechsler Memory Scale



Cognition in family members

Explanation of technical terms

* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small.¹⁵

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified (100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. Less than 0.4 represents a small effect, around 0.5 a medium effect, and over 0.8 represents a large effect.¹⁵

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or < 0.2 ¹⁶. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios



Cognition in family members

measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg, r) indicate the strength of association or relationship between variables. They can provide an indirect indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (b) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent considerable heterogeneity and over this is considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula¹⁵;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, these criteria should be relaxed.¹⁷

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and/or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available and is therefore inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.



Cognition in family members

References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Dickinson D, Ramsey ME, Gold JM (2007): Overlooking the obvious: a meta-analytic comparison of digit symbol coding tasks and other cognitive measures in schizophrenia. *Archives of General Psychiatry* 64: 532-42.
4. Sitskoorn MM, Aleman A, Ebisch SJ, Appels MC, Kahn RS (2004): Cognitive deficits in relatives of patients with schizophrenia: a meta-analysis. *Schizophrenia Research* 71: 285-95.
5. Snitz BE, Macdonald AW, 3rd, Carter CS (2006): Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: a meta-analytic review of putative endophenotypes. *Schizophrenia Bulletin* 32: 179-94.
6. Szoke A, Schurhoff F, Mathieu F, Meary A, Ionescu S, Leboyer M (2005): Tests of executive functions in first-degree relatives of schizophrenic patients: a meta-analysis. *Psychological Medicine* 35: 771-82.
7. Trandafir A, Meary A, Schurhoff F, Leboyer M, Szoke A (2006): Memory tests in first-degree adult relatives of schizophrenic patients: a meta-analysis. *Schizophrenia Research* 81: 217-26.
8. Whyte MC, McIntosh AM, Johnstone EC, Lawrie SM (2005): Declarative memory in unaffected adult relatives of patients with schizophrenia: a systematic review and meta-analysis. *Schizophrenia Research* 78: 13-26.
9. Cohen AS, Brown LA, Auster TL (2012): Olfaction, "olfiction," and the schizophrenia-spectrum: An updated meta-analysis on identification and acuity. *Schizophrenia Research* 135: 152-7.
10. Jameson KG, Nasrallah HA, Northern TG, Welge JA (2011): Executive function impairment in first-degree relatives of persons with schizophrenia: A meta-analysis of controlled studies. *Asian Journal of Psychiatry* 4: 96-9.
11. Bora E, Pantelis C (2013): Theory of mind impairments in first-episode psychosis, individuals at ultra-high risk for psychosis and in first-degree relatives of schizophrenia: Systematic review and meta-analysis. *Schizophrenia Research* doi.org/10.1016/j.schres.2012.12.013.
12. Bora E (2017): A comparative meta-analysis of neurocognition in first-degree relatives of patients with schizophrenia and bipolar disorder. *European Psychiatry: the Journal of the Association of European Psychiatrists* 45: 121-8.
13. Lin SZ, Wu YK, Su YA, Si TM (2019): Prospective memory in non-psychotic first-degree relatives of patients with schizophrenia: A meta-analysis. *Neuropsychiatric Disease and Treatment* 15: 1563-71.
14. Martin D, Croft J, Pitt A, Strelchuk D, Sullivan S, Zammit S (2020): Systematic review and meta-analysis of the relationship between genetic risk for schizophrenia and facial emotion recognition. *Schizophrenia Research* 218: 7-13.
15. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
16. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
17. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. Version 3.2 for Windows