



## **BIPOLAR DISORDERS Factsheet**

**April 2019** 

### What is internalised stigma?

Internalised stigma occurs within an individual, such that a person's attitude may reinforce a negative self-perception of mental disorders, resulting in reduced sense of self-worth, anticipation of social rejection, and a desire for social distance. Stigma can be an important barrier for people with bipolar disorder to seek out proper treatment.

#### What is the evidence for treatments for internalised stigma?

Moderate to high quality evidence suggests a small effect of reduced internalised stigma with targeted interventions comprising psychoeducation and peer-led group discussion.



NeuRA (Neuroscience Research Australia) is one of the largest independent medical and clinical research institutes in Australia and an international leader in neurological research.

Diseases of the brain and nervous system pose the greatest health, economic and social burden of any disease group because they are chronic, debilitating and have no known cures.

Medical research is the cornerstone of efforts to advance the health and wellbeing of families and the community. Our dedicated scientists are focussed on transforming their research into significant and practical benefits for all patients.

While we hope you find this information useful, it is always important to discuss any questions about schizophrenia or its treatment with your doctor or other health care provider.

#### For more information see the technical table

# HOW YOUR SUPPORT HELPS

We are able to make significant advances due to the generosity of countless people. Your donation allows us to continue to work towards transforming lives. For information on how you can support our research, phone 1300 888 019 or make a secure donation at neura.edu.au/donate/schizophrenia.