

For PTSD symptoms

Introduction

Eye Movement Desensitisation and Reprocessing (EMDR) therapy and other trauma-focused therapies are generally beneficial for people with post-traumatic stress disorder (PTSD). They are also currently being investigated in patients with other diagnoses who have had exposure to traumatic events.

EMDR therapy involves identifying past events that are causing dysfunction, identifying current circumstances that elicit distress, and desensitising internal and external triggers. During EMDR therapy, patients attend to the emotionally disturbing material in brief sequential doses while simultaneously focusing on an external stimulus, usually therapist-directed lateral eye movements. The therapist may teach the patient a variety of imagery and stress reduction techniques to use during and between sessions. When patients report no distress related to the targeted memory, they are asked to focus on a preferred positive belief instead. Over the course of treatment, patients gain insight about their stressful situations, and the associated emotional distress reduces.

Other trauma-focused therapies include prolonged exposure, which involves cognitive behavioral therapy to teach people to gradually face trauma-related memories, feelings and situations.

Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2000 that report results separately for people with a diagnosis of schizophrenia or related disorders. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. When multiple copies of review topics were found, only the most recent and/or comprehensive version was included. Reviews with pooled data are prioritised for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist that describes a preferred way to present a meta-analysis¹. Reviews reporting less than 50% of items have been excluded from the library. The PRISMA flow diagram is a suggested way of providing information about studies included and excluded with reasons for exclusion. Where no flow diagram has been presented by individual reviews, but identified studies have been described in the text, reviews have been checked for this item. Note that early reviews may have been guided by less stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons, imprecise or sparse data and high probability of reporting bias. It may also be downgraded if risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large or if there is a dose dependent response. We have also taken into account sample size and whether results are consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)². The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found two reviews that met our inclusion criteria^{3,4}.

For PTSD symptoms

- Moderate to low quality evidence suggests trauma-focused therapies may have long-term effects on delusions and PTSD symptoms and can improve positive symptoms immediately post-treatment. Exposure-based therapies (prolonged exposure, written exposure, elements of imaginal exposure, and EMDR) may be more effective than non-exposure based therapies (cognitive restructuring interventions).

For PTSD symptoms

Brand RM, McEnery C, Rossell S, Bendall S, Thomas N

Do trauma-focussed psychological interventions have an effect on psychotic symptoms? A systematic review and meta-analysis

Schizophrenia Research 2018; 195: 13-22

[View review abstract online](#)

<p>Comparison</p>	<p>Various trauma-focused therapies vs. various control conditions.</p>
<p>Summary of evidence</p>	<p>Moderate to low quality evidence (medium-large samples, inconsistent, precise, indirect) suggests trauma-focused therapies may improve positive symptoms immediately post-treatment, and may have longer-term effects on delusions and PTSD symptoms. Exposure-based therapies (prolonged exposure, written exposure, elements of imaginal exposure, and EMDR) may be more effective than non-exposure based therapies (cognitive restructuring interventions).</p>
<p style="text-align: center;">Symptoms</p>	
<p><i>Trauma-focused therapies had a small, significant effect of improving positive symptoms immediately post-treatment but not at follow-up;</i></p> <p style="padding-left: 40px;">Post-treatment: 5 RCTs, N = 345, $g = 0.31$, 95%CI 0.55 to 0.06, $p = 0.014$</p> <p style="padding-left: 40px;">Follow-up: 5 RCTs, N = 345, $g = 0.18$, 95%CI 0.42 to -0.06, $p = 0.148$</p> <p><i>Trauma-focused therapies had a small, significant effect of improving delusions at follow-up but not immediately post-treatment, although the effect sizes were similar;</i></p> <p style="padding-left: 40px;">Post-treatment: 2 RCTs, N = 216, $g = 0.37$, 95%CI 0.87 to -0.12, $p = 0.139$</p> <p style="padding-left: 40px;">Follow-up: 2 RCTs, N = 216, $g = 0.38$, 95%CI 0.67 to 0.10, $p = 0.008$</p> <p><i>Trauma-focused therapies had a small, significant effect of improving PTSD symptoms at follow-up but not immediately post-treatment;</i></p> <p style="padding-left: 40px;">Post-treatment: 5 RCTs, N = 322, $g = 0.21$, 95%CI 0.70 to -0.27, $p = 0.388$</p> <p style="padding-left: 40px;">Follow-up: 5 RCTs, N = 322, $g = 0.31$, 95%CI 0.62 to 0.00, $p = 0.049$</p> <p>There were no significant effects for hallucinations, negative symptoms, depression, or anxiety. Subgroup analyses found that exposure-based therapies (prolonged exposure, written exposure, elements of imaginal exposure, and EMDR) showing larger effects than non-exposure based therapies (cognitive restructuring interventions) for PTSD symptoms, and to a lesser extent, positive symptoms, hallucinations, and negative symptoms. Longer treatments were also associated with larger effect sizes.</p>	

For PTSD symptoms

Consistency in results[†]	Authors reported moderate to high heterogeneity.
Precision in results[§]	Precise
Directness of results	Indirect; various interventions and control conditions combined.

Valiente-Gomez A, Moreno-Alcazar A, Treen D, Cedron C, Colom F, Perez V, Amann BL

EMDR beyond PTSD: A systematic literature review

Frontiers in Psychology 2017; 8: 1668

[View review abstract online](#)

Comparison	EMDR vs. various control conditions in people with psychotic disorders.
Summary of evidence	Moderate to low quality evidence (small to medium-sized samples, unable to assess consistency or precision, direct) finds some improvements in PTSD symptoms with EMDR treatment, although EMDR may not be any more effective than prolonged exposure treatment.
Symptoms	
<p>1 RCT (N = 155) compared EMDR with prolonged exposure or waitlist and found both trauma-focused treatments were effective at treating PTSD symptoms. Paranoid thoughts improved more with EMDR, while depressive symptoms improved more with prolonged exposure.</p> <p>1 pilot (N = 10) compared EMDR with prolonged exposure treatments or waitlist and found both trauma-focused treatments were effective at treating PTSD symptoms.</p> <p>1 pilot (N = 45) compared EMDR with progressive relaxation and found no improvements within groups, except of an advantage of EMDR for negative symptoms.</p>	
Consistency in results	Unable to assess; no measure of consistency is reported
Precision in results	Unable to assess; no confidence intervals are reported.
Directness of results	Direct

For PTSD symptoms

Explanation of acronyms

CI = confidence interval, g = Hedges' g standardised mean difference, EMDR = Eye Movement Desensitisation and Reprocessing, N = number of participants, p = statistical probability of obtaining that result ($p < 0.05$ generally regarded as significant), PTSD = post-traumatic stress disorder, RCT = randomised controlled trial, vs. = versus

For PTSD symptoms

Explanation of technical terms

* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small⁵.

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion

of actual positives that are correctly identified (100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) which allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. Less than 0.4 represents a small effect, around 0.5 a medium effect, and over 0.8 represents a large effect⁵.

Relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or < 0.2 ⁶. Odds ratios (ORs) are similar to RRs, but they are based on the probability of an event occurring divided by the probability of that event not occurring. ORs and RRs are similar in size when the event is rare, such as with schizophrenia. InOR stands for logarithmic

For PTSD symptoms

OR where a lnOR of 0 shows no difference between groups. Hazard ratios (HRs) measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg, r) indicate the strength of association or relationship between variables. They can provide an indirect indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (b) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent considerable heterogeneity and over this is considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula⁵;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, these criteria should be relaxed⁷.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and/or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available and is therefore inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.

For PTSD symptoms

References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Valiente-Gomez A, Moreno-Alcazar A, Treen D, Cedron C, Colom F, Perez V, *et al.* (2017): EMDR beyond PTSD: A systematic literature review. *Frontiers in Psychology* 8: 1668.
4. Brand RM, McEnery C, Rossell S, Bendall S, Thomas N (2018): Do trauma-focussed psychological interventions have an effect on psychotic symptoms? A systematic review and meta-analysis. *Schizophrenia Research* 195: 13-22.
5. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
6. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
7. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. *Version 3.2 for Windows*.