



## Post-traumatic growth

### Introduction

Post-traumatic growth refers to positive psychological changes that occur after exposure to a traumatic event that are beyond previous levels of functioning. Growth can involve spiritual change, better relationships, more appreciation of life, new possibilities, and increased personal strength. This growth involves both the struggle and symptoms associated with trauma exposure and any positive outcomes that represent adaptation to the traumatic event.

### Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2010 that report results separately for people with PTSD. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. When multiple copies of reviews were found, only the most recent version was included. We prioritised reviews with pooled data for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist that describes a preferred way to present a meta-analysis<sup>1</sup>. Reviews with less than 50% of items checked have been excluded from the library. Note that early reviews may have been guided by less stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons, imprecise or sparse data and high probability of reporting bias. It may also be downgraded if

risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large or if there is a dose dependent response. We have also taken into account sample size and whether results are consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)<sup>2</sup>. The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

### Results

We found two systematic reviews that met our inclusion criteria<sup>3, 4</sup>.

- Moderate to high quality evidence found a small to medium-sized association between increased severity of PTSD symptoms and increased post-traumatic growth. This association was strongest in children, in caregivers, and in people exposed to military conflicts, accidents, or natural disasters. It was lower in people exposed to diseases or those in helping professions and was not present in people exposed to sexual abuse.



Post-traumatic growth

Liu AN, Wang LL, Li HP, Gong J, Liu XH

**Correlation Between Posttraumatic Growth and Posttraumatic Stress Disorder Symptoms Based on Pearson Correlation Coefficient: A Meta-Analysis**

Journal of Nervous and Mental Disease 2017; 205: 380-9

[View review abstract online](#)

<b>Comparison</b>	<b>Association between PTSD symptoms and post-traumatic growth.</b>
<b>Summary of evidence</b>	<b>Moderate to high quality evidence (large sample, inconsistent, precise, direct) found a small to medium-sized association between increased PTSD symptoms and increased post-traumatic growth. This association was strongest in children, in caregivers, and in people exposed to military conflicts, accidents, or natural disasters. It was lower in people exposed to diseases or in people in helping professions. It was not present in people exposed to sexual abuse.</b>
<b>Post-traumatic growth</b>	
<p><i>A small to medium-sized association was found between increased PTSD symptoms and increased post-traumatic growth;</i></p> <p style="text-align: center;">63 studies, N = 26,951, z = 0.22, 95%CI 0.18 to 0.25, I<sup>2</sup> = 86.5%</p> <p style="text-align: center;">The effect size was larger in children than in adults (z = 0.30 vs. 0.21).</p> <p>Effect sizes were similar in caregivers with PTSD (z = 0.33), in people exposed to military conflicts (z = 0.30), accidents (z = 0.32), and natural disasters (z = 0.22), but it was lower in people exposed to diseases (z = 0.13) and in people in helping professions (z = 0.13). It was not significant in people exposed to sexual abuse (z = 0.05).</p> <p style="text-align: center;">There was no evidence of publication bias.</p>	
<b>Consistency in results<sup>‡</sup></b>	Inconsistent
<b>Precision in results<sup>§</sup></b>	Precise
<b>Directness of results<sup>  </sup></b>	Direct

Marziliano A, Tuman M, Moyer A

**The relationship between post-traumatic stress and post-traumatic growth**



**Post-traumatic growth**

**in cancer patients and survivors: A systematic review and meta-analysis**

Psycho-Oncology 2019; 29: 604-16

[View review abstract online](#)

<b>Comparison</b>	Association between PTSD symptoms and post-traumatic growth in people with cancer.
<b>Summary of evidence</b>	Moderate to high quality evidence (large sample, inconsistent, precise, direct) found a small association between increased PTSD symptoms and increased post-traumatic growth in people with cancer. This association was strongest in people with stage four cancer and in studies that used the Impact of Events Scale to measure PTSD symptoms.
<b>Post-traumatic growth</b>	
<p><i>A small association was found between increased PTSD symptoms and increased post-traumatic growth in people with cancer;</i></p> <p>51 studies, N = 16,305, z = 0.08, 95%CI 0.05 to 0.12, p = 0.005, I<sup>2</sup> = 69%</p> <p>The relationship was strongest in studies of patients with stage four cancer (more advanced), and in studies that used the Impact of Events Scale to assess PTSD symptoms.</p> <p>There were no moderating effects of time since diagnosis or of measure used to assess post-traumatic growth.</p> <p>There was no evidence of publication bias.</p>	
<b>Consistency in results</b>	Inconsistent
<b>Precision in results</b>	Precise
<b>Directness of results</b>	Direct

**Explanation of acronyms**

CI = confidence interval, I<sup>2</sup> = the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance), N = number of participants, p = statistical probability of obtaining that result, z = Fisher's z transformation



## Post-traumatic growth

### Explanation of technical terms

\* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small<sup>5</sup>.

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified

(100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. Less than 0.4 represents a small effect, around 0.5 a medium effect, and over 0.8 represents a large effect<sup>5</sup>.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction ( $< 1$ ) or an increase ( $> 1$ ) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if  $RR > 2$  or  $< 0.5$  and a large effect if  $RR > 5$  or  $< 0.2$ <sup>6</sup>. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg,  $r$ ,  $z$ ) indicate the strength of association or relationship



## Post-traumatic growth

between variables. They can provide an indirect indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An  $r$  of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised ( $b$ ) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate.  $I^2$  is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent considerable heterogeneity and over this is considerable heterogeneity.  $I^2$  can be calculated from  $Q$  (chi-square) for the test of heterogeneity with the following formula<sup>5</sup>;

$$I^2 = \left( \frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence

limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, these criteria should be relaxed<sup>7</sup>.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and/or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available and is therefore inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.



## Post-traumatic growth

### References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Liu AN, Wang LL, Li HP, Gong J, Liu XH (2017): Correlation Between Posttraumatic Growth and Posttraumatic Stress Disorder Symptoms Based on Pearson Correlation Coefficient: A Meta-Analysis. *Journal of Nervous and Mental Disease* 205: 380-9.
4. Marziliano A, Tuman M, Moyer A (2019): The relationship between post-traumatic stress and post-traumatic growth in cancer patients and survivors: A systematic review and meta-analysis. *Psycho-Oncology* 29: 604-16.
5. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
6. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
7. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. *Version 3.2 for Windows*