



## Immune system changes

### Introduction

The immune system is the body's first line of defense and predominantly uses proteins called cytokines that are secreted by immune cells and act to allow cell-to-cell communication. Cytokines include interleukins (IL), interferons (IFN), tumor necrosis factors (TNF), transforming growth factors (TGF), and chemokines. These have influence over many cell types, including T helper lymphocytes (Th cells, or white blood cells) and act to regulate immunological and inflammatory responses to pathogens. They are understood to function as intermediaries between the immune system and the central nervous system (CNS). Alterations of these immune-system mediators could have widespread effects for immune system functioning. C-reactive protein (CRP) is released by the body during inflammation. Increased CRP blood levels are not only suggestive of infection, but also chronic inflammatory conditions, including cardiovascular disease, diabetes, and metabolic dysfunction.

### Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2010 that report results separately for people with PTSD. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. When multiple reviews assessing the same topic was found, only the most recent and comprehensive review was included. We prioritised reviews with pooled data for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist that describes a preferred way to present a meta-analysis<sup>1</sup>. Reviews with less than 50% of items checked have been excluded from the library. Note that early reviews may have been guided by less

stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons, imprecise or sparse data and high probability of reporting bias. It may also be downgraded if risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large or if there is a dose dependent response. We have also taken into account sample size and whether results are consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)<sup>2</sup>. The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

### Results

We found one systematic review that met our inclusion criteria<sup>3</sup>.

- Moderate to high quality evidence found large increases in IFN- $\gamma$ , IL-2, IL-1 $\beta$ , and IL-6, and medium-sized increases in TNF- $\alpha$ , WBC, and CRP in people with PTSD compared to controls. There were no differences in IL-4, IL-8, IL-10, lymphocyte, B lymphocyte, CD3+ T lymphocyte, CD4+ T lymphocyte, CD8+ T lymphocyte, or NK cell. The effect sizes became non-significant in medicated patients for IL-1 $\beta$ , IL-2, and IL-6. The effect for CD8+ T lymphocyte became significant when using only trauma-exposed controls (reduced levels in patients). When using only non-trauma-exposed controls, the



## Immune system changes

effect for IL-1 $\beta$  became non-significant, while the effect for IL-10 became significant (increased levels in patients). In patients without comorbid major depression, the effect sizes for IL-8 and CD4+ T lymphocyte became significant (increased levels in patients).



Immune system changes

Yang JJ, Jiang W

**Immune biomarkers alterations in post-traumatic stress disorder: A systematic review and meta-analysis**

Journal of Affective Disorders 2020; 268: 39-46

[View review abstract online](#)

<b>Comparison</b>	<b>Immune biomarker alterations in people with PTSD vs. controls.</b>
<b>Summary of evidence</b>	<p><b>Moderate to high quality evidence (large samples, some inconsistency and imprecision, direct) found large increases in IFN-<math>\gamma</math>, IL-2, IL-1<math>\beta</math>, and IL-6, and medium-sized increases in TNF-<math>\alpha</math>, WBC, and CRP in people with PTSD compared to controls. There were no differences in IL-4, IL-8, IL-10, lymphocyte, B lymphocyte, CD3+ T lymphocyte, CD4+ T lymphocyte, CD8+ T lymphocyte, or NK cell.</b></p> <p><b>The effect sizes became non-significant in medicated patients for IL-1<math>\beta</math>, IL-2, and IL-6. The effect for CD8+ T lymphocyte became significant when using only trauma-exposed controls (reduced levels in patients). When using only non-trauma-exposed controls, the effect for IL-1<math>\beta</math> became non-significant, while the effect for IL-10 became significant (increased levels in patients). In patients without comorbid major depression, the effect sizes for IL-8 and CD4+ T lymphocyte became significant (increased levels in patients).</b></p>
<b>Immune biomarkers</b>	
<p><i>The following biomarkers were increased in people with PTSD vs. controls (in order of descending effect size);</i></p> <p>IFN-<math>\gamma</math>: 5 studies, N = 441, SMD = 1.68, 95%CI 0.54 to 2.82, <math>p = 0.004</math>, <math>I^2 = 96\%</math>              IL-2: 7 studies, N = 475, SMD = 1.53, 95%CI 0.45 to 2.61, <math>p = 0.006</math>, <math>I^2 = 96\%</math>              IL-1<math>\beta</math>: 9 studies, N = 669, SMD = 0.98, 95%CI 0.20 to 1.77, <math>p = 0.01</math>, <math>I^2 = 95\%</math>              IL-6: 20 studies, N = 1,139, SMD = 0.89, 95%CI 0.42 to 1.36, <math>p = 0.0002</math>, <math>I^2 = 92\%</math>              TNF-<math>\alpha</math>: 14 studies, N = 1,219, SMD = 0.67, 95%CI 0.21 to 1.13, <math>p = 0.004</math>, <math>I^2 = 92\%</math>              WBC: 4 studies, N = 998, SMD = 0.29, 95%CI 0.07 to 0.52, <math>p = 0.01</math>, <math>I^2 = 45\%</math>              CRP: 15 studies, N = 4,672, SMD = 0.27, 95%CI 0.12 to 0.42, <math>p = 0.0003</math>, <math>I^2 = 45\%</math></p> <p>There were no significant differences in IL-4, IL-8, IL-10, lymphocyte, B lymphocyte, CD3+ T lymphocyte, CD4+ T lymphocyte, CD8+ T lymphocyte, or NK cell.</p> <p>Subgroup analyses showed the effect sizes became non-significant in medicated patients for IL-1<math>\beta</math>, IL-2, and IL-6.</p>	



**Immune system changes**

<p>The effect for CD8+ T lymphocyte became significant when using only trauma-exposed controls (reduced levels in patients).</p> <p>When using only non-trauma-exposed controls, the effect for IL-1<math>\beta</math> became non-significant, while the effect for IL-10 became significant (increased levels in patients).</p> <p>In patients without comorbid major depression, the effect sizes for IL-8 and CD4+ T lymphocyte became significant (increased levels in patients).</p>	
<b>Consistency in results<sup>‡</sup></b>	Inconsistent, apart from WBC and CRP
<b>Precision in results<sup>§</sup></b>	Imprecise, apart from WBC and CRP
<b>Directness of results<sup>  </sup></b>	Direct

**Explanation of acronyms**

CI = confidence interval, I<sup>2</sup> = the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance), N = number of participants, p = statistical probability of obtaining that result, SMD = standardised mean difference, vs. = versus



## Immune system changes

### Explanation of technical terms

\* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small<sup>4</sup>.

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified

(100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. Less than 0.4 represents a small effect, around 0.5 a medium effect, and over 0.8 represents a large effect<sup>4</sup>.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction ( $< 1$ ) or an increase ( $> 1$ ) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if  $RR > 2$  or  $< 0.5$  and a large effect if  $RR > 5$  or  $< 0.2$ <sup>5</sup>. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg,  $r$ ) indicate the strength of association or relationship



## Immune system changes

between variables. They can provide an indirect indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An  $r$  of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised ( $b$ ) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate.  $I^2$  is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent considerable heterogeneity and over this is considerable heterogeneity.  $I^2$  can be calculated from  $Q$  (chi-square) for the test of heterogeneity with the following formula<sup>4</sup>;

$$I^2 = \left( \frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence

limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, these criteria should be relaxed<sup>6</sup>.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and/or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available and is therefore inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.



## Immune system changes

### References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Yang JJ, Jiang W (2020): Immune biomarkers alterations in post-traumatic stress disorder: A systematic review and meta-analysis. *Journal of Affective Disorders* 268: 39-46.
4. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
5. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
6. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. *Version 3.2 for Windows*