



## Prevalence in elderly people

### Introduction

Prevalence represents the overall proportion of individuals in a population who have the disorder of interest. It is different from incidence, which represents only the new cases that have developed over a particular time period. Point prevalence is the proportion of individuals in a population who have the disorder at a given point in time (e.g., at one-month post-trauma), while period prevalence is the proportion of individuals in a population who have the disorder over specific time periods (e.g., one to two months post-trauma). Lifetime prevalence is the proportion of individuals in a population who have ever had the disorder and lifetime morbid risk also includes those who had the disorder but were deceased at the time of the survey. This summary table presents the evidence on prevalence rates of PTSD in elderly people exposed to trauma.

### Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2010 that report results separately for people with PTSD. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. When multiple copies of reviews were found, only the most recent version was included. We prioritised reviews with pooled data for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist that describes a preferred way to present a meta-analysis<sup>1</sup>. Reviews with less than 50% of items checked have been excluded from the library. Note that early reviews may have been guided by less stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development

and Evaluation ([GRADE](#)) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons, imprecise or sparse data and high probability of reporting bias. It may also be downgraded if risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large or if there is a dose dependent response. We have also taken into account sample size and whether results are consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)<sup>2</sup>. The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

### Results

We found four systematic reviews that met our inclusion criteria<sup>3-6</sup>.

- Moderate quality evidence finds the 12-month prevalence rate of PTSD in older adults ( $\geq 65$  years) is 1.56%, and the lifetime prevalence rate is 2.66%. Lifetime rates of PTSD are nearly twice as high in women than in men.
- Moderate quality evidence finds the average prevalence of PTSD in older people ( $\geq 65$  years) after a fall is 27.5%, which represents a small, significant increase in risk compared to older people with no previous fall.
- Moderate quality evidence finds the prevalence rate of PTSD in older US veterans ( $\geq 65$  years) is 8.4%.
- Moderate to low quality evidence finds the prevalence rate of PTSD in older prisoners ( $\geq 50$  years) is 6.2%.



Prevalence in elderly people

*Bloch F*

**Literature review and meta-analysis of risk factors for delayed post-traumatic stress disorder in older adults after a fall**

International Journal of Geriatric Psychiatry 2017; 32: 136-40

[View review abstract online](#)

<b>Comparison</b>	Prevalence of PTSD in older adults (≥65 years) after a fall (up to 24 weeks).
<b>Summary of evidence</b>	Moderate quality evidence (small to medium-sized samples, consistent, imprecise, direct) finds the average prevalence of PTSD after a fall is 27.5%, which represents a small, significant increase in risk compared to older people with no previous fall.
<b>Prevalence in older adults after a fall</b>	
3 studies, N = 211, prevalence = 27.5%	
<i>A small, significant increase in the risk of PTSD in older people with a fall compared to older people without a fall;</i>	
2 studies, OR = 2.79, 95%CI 1.03 to 7.53, <i>p</i> < 0.05	
<b>Consistency in results</b>	Authors report the OR data are consistent.
<b>Precision in results</b>	Imprecise for OR.
<b>Directness of results</b>	Direct

*Di Lorito C, Vollm B, Dening T*

**Psychiatric disorders among older prisoners: a systematic review and comparison study against older people in the community**

Aging & Mental Health 2018; 22: 1-10

[View review abstract online](#)

<b>Comparison</b>	Prevalence of PTSD in older prisoners (≥50 years).
<b>Summary of evidence</b>	Moderate to low quality evidence (unclear sample size, direct) finds the prevalence of PTSD older prisoners is 6.2%.



**Prevalence in elderly people**

<b>Prevalence in older prisoners</b>	
3 studies, N not reported, point prevalence = 6.2%	
<b>Consistency in results</b>	Unable to assess; no measure of consistency is reported.
<b>Precision in results</b>	Unable to assess; no measure of precision is reported.
<b>Directness of results</b>	Direct

<p><i>Grenier S, Payette MC, Gunther B, Askari S, Desjardins FF, Raymond B, Berbiche D</i></p> <p><b>Association of age and gender with anxiety disorders in older adults: A systematic review and meta-analysis</b></p> <p>International Journal of Geriatric Psychiatry 2019; 34: 397-407</p> <p><a href="#">View review abstract online</a></p>	
<b>Comparison</b>	<b>Prevalence of PTSD in older adults (≥65 years).</b>
<b>Summary of evidence</b>	<b>Moderate quality evidence (large samples, inconsistent, imprecise, direct) finds the 12-month prevalence rate of PTSD in older adults (&gt;65 years) is 1.56%, and the lifetime prevalence rate is 2.66%. Lifetime rates are nearly twice as high in women than in men.</b>
<b>Prevalence in older adults</b>	
<p>5 studies, N = 15,177, 12-month prevalence = 1.56%, 95%CI 0.98% to 2.47%, I<sup>2</sup> = 88.8%</p> <p>5 studies, N = 11,746, lifetime prevalence = 2.66%, 95%CI 2.19% to 2.33%, I<sup>2</sup> = 50.6%</p> <p><i>Lifetime prevalence was higher in women than in men;</i></p> <p>Lifetime prevalence = 3.42% vs. 1.83%, OR = 1.93, 95%CI 1.24 to 3.03, p = 0.002, I<sup>2</sup> = 0%</p>	
<b>Consistency in results</b>	Inconsistent
<b>Precision in results</b>	Imprecise
<b>Directness of results</b>	Direct

*Williamson V, Stevelink SAM, Greenberg K, Greenberg N*



Prevalence in elderly people

**Prevalence of Mental Health Disorders in Elderly U.S. Military Veterans: A Meta-Analysis and Systematic Review**

American Journal of Geriatric Psychiatry 2018; 26: 534-45

[View review abstract online](#)

<b>Comparison</b>	Prevalence of PTSD in older US veterans (≥65 years).
<b>Summary of evidence</b>	Moderate quality evidence (large sample, inconsistent, imprecise, direct) finds the prevalence of PTSD in older US veterans (≥65 years) is 8.4%.
<b>Prevalence in older US veterans</b>	
8 studies, N = 1,296,967, prevalence = 8.4%, 95%CI 2.04% to 17.88%, I <sup>2</sup> = 99.99%	
<b>Consistency in results</b>	Inconsistent
<b>Precision in results</b>	Appears imprecise
<b>Directness of results</b>	Direct

Explanation of acronyms

CI = confidence interval, I<sup>2</sup> = the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance), N = number of participants, OR = odds ratio, vs. = versus



## Prevalence in elderly people

### Explanation of technical terms

\* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small<sup>7</sup>.

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified

(100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. Less than 0.4 represents a small effect, around 0.5 a medium effect, and over 0.8 represents a large effect<sup>7</sup>.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction ( $< 1$ ) or an increase ( $> 1$ ) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if  $RR > 2$  or  $< 0.5$  and a large effect if  $RR > 5$  or  $< 0.2$ <sup>8</sup>. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg,  $r$ ) indicate the strength of association or relationship





## Prevalence in elderly people

between variables. They can provide an indirect indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An  $r$  of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised ( $b$ ) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

‡ Inconsistency refers to differing estimates of effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate.  $I^2$  is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent considerable heterogeneity and over this is considerable heterogeneity.  $I^2$  can be calculated from  $Q$  (chi-square) for the test of heterogeneity with the following formula<sup>7</sup>;

$$I^2 = \left( \frac{Q - df}{Q} \right) \times 100\%$$

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence

limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, these criteria should be relaxed<sup>9</sup>.

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and/or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available and is therefore inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.



## Prevalence in elderly people

### References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Bloch F (2017): Literature review and meta-analysis of risk factors for delayed post-traumatic stress disorder in older adults after a fall. *International Journal of Geriatric Psychiatry* 32: 136-40.
4. Di Lorito C, Vollm B, Denning T (2018): Psychiatric disorders among older prisoners: a systematic review and comparison study against older people in the community. *Aging & Mental Health* 22: 1-10.
5. Grenier S, Payette MC, Gunther B, Askari S, Desjardins FF, Raymond B, *et al.* (2019): Association of age and gender with anxiety disorders in older adults: A systematic review and meta-analysis. *International Journal of Geriatric Psychiatry* 34: 397-407.
6. Williamson V, Stevelink SAM, Greenberg K, Greenberg N (2018): Prevalence of Mental Health Disorders in Elderly U.S. Military Veterans: A Meta-Analysis and Systematic Review. *American Journal of Geriatric Psychiatry* 26: 534-45.
7. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
8. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
9. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. *Version 3.2 for Windows*