



Prevalence in criminal offenders

Introduction

Many studies have reported a high prevalence of various health problems, including mental health problems, among people in forensic settings. Prevalence represents the overall proportion of individuals in a population who have the disorder of interest. It is different from incidence, which represents only the new cases that have developed over a particular time-period. Point prevalence is the proportion of individuals in a population who have the disorder at a given point in time (e.g., at one-month post-trauma), while period prevalence is the proportion of individuals in a population who have the disorder over specific time periods (e.g., one to two months post-trauma). Lifetime prevalence is the proportion of individuals in a population who have ever had the disorder and lifetime morbid risk also includes those who had the disorder but were deceased at the time of the survey.

Method

We have included only systematic reviews (systematic literature search, detailed methodology with inclusion/exclusion criteria) published in full text, in English, from the year 2010 that report results separately for people with PTSD. Reviews were identified by searching the databases MEDLINE, EMBASE, and PsycINFO. When multiple copies of reviews were found, only the most recent version was included. We prioritised reviews with pooled data for inclusion.

Review reporting assessment was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist that describes a preferred way to present a meta-analysis¹. Reviews with less than 50% of items checked have been excluded from the library. Note that early reviews may have been guided by less stringent reporting checklists than the PRISMA, and that some reviews may have been limited by journal guidelines.

Evidence was graded using the Grading of Recommendations Assessment, Development and Evaluation ([GRADE](#)) Working Group approach where high quality evidence such as that gained from randomised controlled trials (RCTs) may be downgraded to moderate or low if review and study quality is limited, if there is inconsistency in results, indirect comparisons, imprecise or sparse data and high probability of reporting bias. It may also be downgraded if risks associated with the intervention or other matter under review are high. Conversely, low quality evidence such as that gained from observational studies may be upgraded if effect sizes are large or if there is a dose dependent response. We have also taken into account sample size and whether results are consistent, precise and direct with low associated risks (see end of table for an explanation of these terms)². The resulting table represents an objective summary of the available evidence, although the conclusions are solely the opinion of staff of NeuRA (Neuroscience Research Australia).

Results

We found three systematic reviews that met our inclusion criteria³⁻⁵.

- Moderate quality evidence finds the point prevalence in male prisoners is 6.2%, and in female prisoners it is 21.1%. One-year prevalence in male prisoners is 9.9%, and in female prisoners it is 26.1%. Lifetime prevalence in male prisoners is 17.8%, and in female prisoners it is 40.4%. These rates were significantly higher in females and were also higher in studies from high-income countries.
- Moderate to low quality evidence finds the prevalence of PTSD older prisoners (>50 years) is 6.2%.
- Moderate quality evidence finds the prevalence of PTSD in adolescent males in detention centres is 8.6% and 18.2% in adolescent females in detention centres.



Prevalence in criminal offenders

Baranyi G, Cassidy M, Fazel S, Priebe S, Mundt AP

Prevalence of Posttraumatic Stress Disorder in Prisoners

Epidemiologic Reviews 2018; 40: 134-45

[View review abstract online](#)

Comparison	Prevalence of PTSD in prisoners.
Summary of evidence	Moderate quality evidence (large samples, inconsistent, imprecise, direct) finds the point prevalence in male prisoners is 6.2%, and in female prisoners it is 21.1%. One-year prevalence in male prisoners is 9.9%, and in female prisoners it is 26.1%. Lifetime prevalence in male prisoners is 17.8%, and in female prisoners it is 40.4%. These rates were significantly higher for females, and also higher in studies from high-income countries.
Prevalence in prisoners	
<p>Point prevalence reported in 50 samples, N = 19,011 from 20 countries</p> <p style="padding-left: 40px;">Males = 6.2%, 95%CI 3.9% to 9.0%, I² = 97%</p> <p style="padding-left: 40px;">Females = 21.1%, 95%CI 16.9% to 25.6%, I² = 90%</p> <p>One-year prevalence reported in 12 samples, N = 4,889 from 4 countries</p> <p style="padding-left: 40px;">Males = 9.9%, 95%CI 3.0% to 20.2%, I² = 99%</p> <p style="padding-left: 40px;">Females = 26.1%, 95%CI 15.9% to 37.8%, I² = 96%</p> <p>Lifetime prevalence reported in 23 samples, N = 9,202 from 7 countries</p> <p style="padding-left: 40px;">Males = 17.8%, 95%CI 12.4% to 23.9%, I² = 97%</p> <p style="padding-left: 40px;">Females = 40.4%, 95%CI 31.8% to 49.3%, I² = 96%</p> <p>Significantly higher prevalence was reported in females, and in studies from high-income countries. These variables accounted for most of the between-sample variance.</p>	
Consistency in results	Inconsistent
Precision in results	Appears imprecise
Directness of results	Direct

Beaudry G, Yu R, Langstrom N, Fazel FS

An Updated Systematic Review and Meta-regression Analysis: Mental



Prevalence in criminal offenders

Disorders Among Adolescents in Juvenile Detention and Correctional Facilities

Journal of the American Academy of Child and Adolescent Psychiatry 2021; 60(1): 46-60

[View review abstract online](#)

Comparison	Prevalence of PTSD in adolescents in juvenile detention and correctional facilities.
Summary of evidence	Moderate quality evidence (large sample, inconsistent, imprecise, direct) finds the prevalence of PTSD in adolescent males in detention centres is 8.6% and 18.2% in adolescent females in detention centres.
Prevalence in adolescents in detention	
Males: 20 studies, N = 14,260, prevalence = 8.6%, 95%CI 6.4% to 10.7%, I ² = 92% Females: 11 studies, N = 1,876, prevalence = 18.2%, 95%CI 13.1% to 23.2%, I ² = 78%	
Consistency in results	Inconsistent
Precision in results	Appears imprecise
Directness of results	Direct

Di Lorito C, Vollm B, Dening T

Psychiatric disorders among older prisoners: a systematic review and comparison study against older people in the community

Aging & Mental Health 2018; 22: 1-10

[View review abstract online](#)

Comparison	Prevalence of PTSD in older prisoners (>50 years).
Summary of evidence	Moderate to low quality evidence (unclear sample size, direct) finds the prevalence of PTSD older prisoners is 6.2%.
Prevalence in older prisoners	
3 studies, N not reported, point prevalence = 6.2%	
Consistency in results	Unable to assess; no measure of consistency is reported.
Precision in results	Unable to assess; no measure of precision is reported.



Prevalence in criminal offenders

Directness of results	Direct
------------------------------	--------

Explanation of acronyms

CI = confidence interval, I^2 = the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance), N = number of participants



Prevalence in criminal offenders

Explanation of technical terms

* Bias has the potential to affect reviews of both RCT and observational studies. Forms of bias include; reporting bias – selective reporting of results; publication bias - trials that are not formally published tend to show less effect than published trials, further if there are statistically significant differences between groups in a trial, these trial results tend to get published before those of trials without significant differences; language bias – only including English language reports; funding bias - source of funding for the primary research with selective reporting of results within primary studies; outcome variable selection bias; database bias - including reports from some databases and not others; citation bias - preferential citation of authors. Trials can also be subject to bias when evaluators are not blind to treatment condition and selection bias of participants if trial samples are small⁶.

† Different effect measures are reported by different reviews.

Prevalence refers to how many existing cases there are at a particular point in time. Incidence refers to how many new cases there are per population in a specified time period. Incidence is usually reported as the number of new cases per 100,000 people per year. Alternatively some studies present the number of new cases that have accumulated over several years against a person-years denominator. This denominator is the sum of individual units of time that the persons in the population are at risk of becoming a case. It takes into account the size of the underlying population sample and its age structure over the duration of observation.

Reliability and validity refers to how accurate the instrument is. Sensitivity is the proportion of actual positives that are correctly identified

(100% sensitivity = correct identification of all actual positives) and specificity is the proportion of negatives that are correctly identified (100% specificity = not identifying anyone as positive if they are truly not).

Weighted mean difference scores refer to mean differences between treatment and comparison groups after treatment (or occasionally pre to post treatment) and in a randomised trial there is an assumption that both groups are comparable on this measure prior to treatment. Standardised mean differences are divided by the pooled standard deviation (or the standard deviation of one group when groups are homogenous) that allows results from different scales to be combined and compared. Each study's mean difference is then given a weighting depending on the size of the sample and the variability in the data. Less than 0.4 represents a small effect, around 0.5 a medium effect, and over 0.8 represents a large effect⁶.

Odds ratio (OR) or relative risk (RR) refers to the probability of a reduction (< 1) or an increase (> 1) in a particular outcome in a treatment group, or a group exposed to a risk factor, relative to the comparison group. For example, a RR of 0.75 translates to a reduction in risk of an outcome of 25% relative to those not receiving the treatment or not exposed to the risk factor. Conversely, a RR of 1.25 translates to an increased risk of 25% relative to those not receiving treatment or not having been exposed to a risk factor. A RR or OR of 1.00 means there is no difference between groups. A medium effect is considered if $RR > 2$ or < 0.5 and a large effect if $RR > 5$ or < 0.2 ⁷. InOR stands for logarithmic OR where a InOR of 0 shows no difference between groups. Hazard ratios measure the effect of an explanatory variable on the hazard or risk of an event.

Correlation coefficients (eg, r) indicate the strength of association or relationship



Prevalence in criminal offenders

between variables. They can provide an indirect indication of prediction, but do not confirm causality due to possible and often unforeseen confounding variables. An r of 0.10 represents a weak association, 0.25 a medium association and 0.40 and over represents a strong association. Unstandardised (b) regression coefficients indicate the average change in the dependent variable associated with a 1 unit change in the independent variable, statistically controlling for the other independent variables. Standardised regression coefficients represent the change being in units of standard deviations to allow comparison across different scales.

limit crosses an effect size of 0.5 in either direction, and for binary and correlation data, an effect size of 0.25. GRADE also recommends downgrading the evidence when sample size is smaller than 300 (for binary data) and 400 (for continuous data), although for some topics, these criteria should be relaxed⁸.

‡ Inconsistency refers to differing estimates of effect across studies (i.e. heterogeneity or variability in results) that is not explained by subgroup analyses and therefore reduces confidence in the effect estimate. I^2 is the percentage of the variability in effect estimates that is due to heterogeneity rather than sampling error (chance) - 0% to 40%: heterogeneity might not be important, 30% to 60%: may represent moderate heterogeneity, 50% to 90%: may represent considerable heterogeneity and over this is considerable heterogeneity. I^2 can be calculated from Q (chi-square) for the test of heterogeneity with the following formula⁶;

$$I^2 = \left(\frac{Q - df}{Q} \right) \times 100\%$$

|| Indirectness of comparison occurs when a comparison of intervention A versus B is not available but A was compared with C and B was compared with C that allows indirect comparisons of the magnitude of effect of A versus B. Indirectness of population, comparator and/or outcome can also occur when the available evidence regarding a particular population, intervention, comparator, or outcome is not available and is therefore inferred from available evidence. These inferred treatment effect sizes are of lower quality than those gained from head-to-head comparisons of A and B.

§ Imprecision refers to wide confidence intervals indicating a lack of confidence in the effect estimate. Based on GRADE recommendations, a result for continuous data (standardised mean differences, not weighted mean differences) is considered imprecise if the upper or lower confidence



Prevalence in criminal offenders

References

1. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009): Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *British Medical Journal* 151: 264-9.
2. GRADE Working Group (2004): Grading quality of evidence and strength of recommendations. *British Medical Journal* 328: 1490.
3. Baranyi G, Cassidy M, Fazel S, Priebe S, Mundt AP (2018): Prevalence of Posttraumatic Stress Disorder in Prisoners. *Epidemiologic Reviews* 40: 134-45.
4. Beaudry G, Yu R, Långström N, Fazel S (2021): An Updated Systematic Review and Meta-regression Analysis: Mental Disorders Among Adolescents in Juvenile Detention and Correctional Facilities. *Journal of the American Academy of Child and Adolescent Psychiatry* 60: 46-60.
5. Di Lorito C, Vollm B, Dening T (2018): Psychiatric disorders among older prisoners: a systematic review and comparison study against older people in the community. *Aging & Mental Health* 22: 1-10.
6. Cochrane Collaboration (2008): Cochrane Handbook for Systematic Reviews of Interventions. Accessed 24/06/2011.
7. Rosenthal JA (1996): Qualitative Descriptors of Strength of Association and Effect Size. *Journal of Social Service Research* 21: 37-59.
8. GRADEpro (2008): [Computer program]. Jan Brozek, Andrew Oxman, Holger Schünemann. Version 3.2 for Windows